

Product datasheet for RR202933L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Gtf2i (NM_001001512) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Gtf2i (NM_001001512) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Gtf2i

Synonyms: Gtf2ird1

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001001512

ORF Size: 2739 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR202933).

OTI Disclaimer:

Sequence:

r: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001001512.2</u>, <u>NP 001001512.2</u>

 RefSeq Size:
 3988 bp

 RefSeq ORF:
 2742 bp

 Locus ID:
 353256

 UniProt ID:
 Q5U2Y1

 Cytogenetics:
 12q12







Gene Summary:

Interacts with the basal transcription machinery by coordinating the formation of a multiprotein complex at the C-FOS promoter, and linking specific signal responsive activator complexes. Promotes the formation of stable high-order complexes of SRF and PHOX1 and interacts cooperatively with PHOX1 to promote serum-inducible transcription of a reporter gene deriven by the C-FOS serum response element (SRE). Acts as a coregulator for USF1 by binding independently two promoter elements, a pyrimidine-rich initiator (Inr) and an upstream E-box (By similarity). Required for the formation of functional ARID3A DNA-binding complexes and for activation of immunoglobulin heavy-chain transcription upon B-lymphocyte activation (By similarity). [UniProtKB/Swiss-Prot Function]