

## Product datasheet for **RR203834**

### Chmp5 (NM\_001025410) Rat Tagged ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Chmp5 (NM_001025410) Rat Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Chmp5
Synonyms:	RGD1305968
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
ORF Nucleotide Sequence:	>RR203834 representing NM_001025410 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGAACCGATTCTTCGAAAAGCGAAACCAAGGCTCCGCCACCTAGCCTGACTGACTGCATTGGGACGG  
 TGGATAGCAGGGCAGAATCCATTGACAAAAAGATTTCCCGTTGGATGCTGAACTGGTGAAATATAAGGA  
 CCAATCAAGAAGATGAGAGAGGGTCTGCTAAGAACATGGTCAAACAGAAAGCCCTGAGGGTTTTAAAG  
 CAAAAACGGATGTATGAGCAGCAGCGAGACAACCTTGCCAACAGTCCTTTAACATGGAGCAAGCTAATT  
 ACACCATCCAGTCACTGAAGGATACCAAAACACGGTTGATGCAATGAAATTGGGAGTAAAGGAAATGAA  
 GAAGGCTTACAAGGAAGTAAAAATTGACCAGATTGAGGATTTACAAGACCAGTTGGAAGACATGATGGAA  
 GATGCAATGAGATCCAGGAGGCCCTGGGCCGAGCTACGGCACCCAGAACTGGATGAGGACGACCTGG  
 AAGCAGAGTTGGATGCGCTGGGCGATGAGCTTCTGGCTGATGAGGACAGCTCTTACTTGGACGAAGCAGC  
 TTCTGCACCGCAATTCAGAGGGCGTCCCACTGACACAAAAACAAGGATGGAGTGCTGGTGGATGAA  
 TTTGGACTGCCACAGATTCCAGCTTCG

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)

**Protein Sequence:** >RR203834 representing NM\_001025410  
 Red=Cloning site Green=Tags(s)

MNRFFGKAKPKAPPSL TDCIGTVDSRAESIDKKISRLDAELVKYKDQIKKMREGPAKNMVQKALRVLK  
 QKRMYYEQQRDLAQQSFNMEQANYTIQSLKDTKTTVDAMKLGVKEMKKAYKEVKIDQIEDLQDQLEDMMME  
 DANEIQEALGRSYGTPELDEDDLEAELDALGDELLADESSYLADEAASAPAIPEGVPTDTKNKDGVLVDE  
 FGLPQIPAS



TRTRPLEQKLISEEDLAANDILDYKDDDDKV

**Restriction Sites:**

Sgfl-MluI

**Cloning Scheme:**

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF

**ACCN:** NM\_001025410

**ORF Size:** 657 bp


**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<u>NM_001025410.1, NP_001020581.1</u>
<b>RefSeq Size:</b>	1355 bp
<b>RefSeq ORF:</b>	660 bp
<b>Locus ID:</b>	297995
<b>UniProt ID:</b>	<u>Q4QQV8</u>
<b>Cytogenetics:</b>	5q22
<b>MW:</b>	24.6 kDa
<b>Gene Summary:</b>	<p>Probable peripherally associated component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4 (By similarity).[UniProtKB/Swiss-Prot Function]</p>

## Product images:



Circular map for RR203834