

## Product datasheet for **RR204100L4V**

### Sh3glb1 (NM\_001011929) Rat Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | Sh3glb1 (NM_001011929) Rat Tagged ORF Clone Lentiviral Particle  |
| Symbol:                   | Sh3glb1  |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-mGFP-P2A-Puro (PS100093)  |
| Tag:                      | mGFP   |
| ACCN:                     | NM_001011929   |
| ORF Size:                 | 1095 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RR204100).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_001011929.1</a> , <a href="#">NP_001011929.1</a>  |
| RefSeq Size:              | 1795 bp  |
| RefSeq ORF:               | 1098 bp  |
| Locus ID:                 | 292156   |
| UniProt ID:               | <a href="#">Q6AYE2</a>   |
| Cytogenetics:             | 2q44   |



[View online »](#)

**Gene Summary:**

May be required for normal outer mitochondrial membrane dynamics. Required for coatamer-mediated retrograde transport in certain cells. May recruit other proteins to membranes with high curvature. May promote membrane fusion. Involved in activation of caspase-dependent apoptosis by promoting BAX/BAK1 activation. Involved in caspase-independent apoptosis during nutrition starvation and involved in the regulation of autophagy. Activates lipid kinase activity of PIK3C3 during autophagy probably by associating with the PI3K complex II (PI3KC3-C2). Associated with PI3KC3-C2 during autophagy may regulate the trafficking of ATG9A from the Golgi complex to the peripheral cytoplasm for the formation of autophagosomes by inducing Golgi membrane tubulation and fragmentation. Involved in regulation of degradative endocytic trafficking and cytokinesis, probably in the context of PI3KC3-C2 (By similarity).[UniProtKB/Swiss-Prot Function]