

## Product datasheet for RR204120L4V

## Eri1 (NM\_001014143) Rat Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Eri1 (NM_001014143) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Eri1
Synonyms:	RGD1308378; Thex1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001014143
ORF Size:	1035 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR204120).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001014143.1, NP 001014165.1</u>
RefSeq Size:	2896 bp
RefSeq ORF:	1038 bp
Locus ID:	361159
UniProt ID:	Q5FVR4
Cytogenetics:	16q12.2



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



Gene Summary:RNA exonuclease that binds to the 3'-end of histone mRNAs and degrades them, suggesting<br/>that it plays an essential role in histone mRNA decay after replication. A 2' and 3'-hydroxyl<br/>groups at the last nucleotide of the histone 3'-end is required for efficient degradation of RNA<br/>substrates. Also able to degrade the 3'-overhangs of short interfering RNAs (siRNAs) in vitro,<br/>suggesting a possible role as regulator of RNA interference (RNAi). Binds with high affinity to<br/>the 3' side of the stem-loop structure and to the downstream cleavage product (DCP) of<br/>histone pre-mRNAs. Requires for binding the 5'-ACCCA-3' sequence present in stem-loop<br/>structure. Able to bind other mRNAs. Required for 5.8S rRNA 3'-end processing. Also binds to<br/>5.8s ribosomal RNA (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US