

## Product datasheet for RR204503L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Trip13 (NM 001011930) Rat Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Trip13 (NM\_001011930) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Trip13

**Mammalian Cell** Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Myc-DDK Tag:

ACCN: NM\_001011930

**ORF Size:** 1296 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RR204503).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

This clone was engineered to express the complete ORF with an expression tag. Expression **OTI Annotation:** 

varies depending on the nature of the gene.

RefSeq: NM 001011930.1, NP 001011930.1

RefSeq Size: 1821 bp RefSeq ORF: 1299 bp Locus ID: 292206 **UniProt ID:** Q5XHZ9

Cytogenetics: 1p11







## **Gene Summary:**

Plays a key role in chromosome recombination and chromosome structure development during meiosis. Required at early steps in meiotic recombination that leads to non-crossovers pathways. Also needed for efficient completion of homologous synapsis by influencing crossover distribution along the chromosomes affecting both crossovers and non-crossovers pathways. Also required for development of higher-order chromosome structures and is needed for synaptonemal-complex formation. In males, required for efficient synapsis of the sex chromosomes and for sex body formation. Promotes early steps of the DNA double-strand breaks (DSBs) repair process upstream of the assembly of RAD51 complexes. Required for depletion of HORMAD1 and HORMAD2 from synapsed chromosomes. Plays a role in mitotic spindle assembly checkpoint (SAC) activation (By similarity).[UniProtKB/Swiss-Prot Function]