

Product datasheet for **RR204754L4V**

Rac1 (NM_134366) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Rac1 (NM_134366) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Rac1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_134366
ORF Size:	576 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR204754).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_134366.1 , NP_599193.1
RefSeq Size:	792 bp
RefSeq ORF:	579 bp
Locus ID:	363875
UniProt ID:	Q6RUV5
Cytogenetics:	12p11



[View online »](#)

Gene Summary:

Plasma membrane-associated small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate cellular responses such as secretory processes, phagocytosis of apoptotic cells, epithelial cell polarization, neurons adhesion, migration and differentiation, and growth-factor induced formation of membrane ruffles (PubMed:16040606, PubMed:16549782). Rac1 p21/rho GDI heterodimer is the active component of the cytosolic factor sigma 1, which is involved in stimulation of the NADPH oxidase activity in macrophages. Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly. Stimulates PKN2 kinase activity (By similarity). In concert with RAB7A, plays a role in regulating the formation of RBs (ruffled borders) in osteoclasts (PubMed:16040606). In glioma cells, promotes cell migration and invasion (PubMed:20696765). In podocytes, promotes nuclear shuttling of NR3C2; this modulation is required for a proper kidney functioning (PubMed:19029984). Required for atypical chemokine receptor ACKR2-induced LIMK1-PAK1-dependent phosphorylation of cofilin (CFL1) and for up-regulation of ACKR2 from endosomal compartment to cell membrane, increasing its efficiency in chemokine uptake and degradation (By similarity). In neurons, is involved in dendritic spine formation and synaptic plasticity (PubMed:25498153). In synapses, seems to mediate the regulation of F-actin cluster formation performed by SHANK3 (PubMed:24089484).[UniProtKB/Swiss-Prot Function]