

Product datasheet for RR204917

Pnrc2 (NM_001103360) Rat Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Pnrc2 (NM_001103360) Rat Tagged ORF Clone

Tag: Myc-DDK

Symbol: Pnrc2

Synonyms: MGC93828

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RR204917 representing NM_001103360
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RR204917 representing NM_001103360

Red=Cloning site Green=Tags(s)

MGGGERYNIPDPQSRNASKNQQQHNRQKTKDQNSQMKIVHKKKERGHGYNPSAVQNGGKTKSLSNNSNWN

ASLSSPSLLFKSQASQNYAGAKFSEPPSPSVLPKPPSHWVHVSLNPSDKETMTFQLKTLLKVQV

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001103360

ORF Size: 402 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

Pnrc2 (NM_001103360) Rat Tagged ORF Clone - RR204917

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001103360.1</u>, <u>NP 001096830.1</u>

 RefSeq Size:
 1899 bp

 RefSeq ORF:
 405 bp

 Locus ID:
 100125373

 UniProt ID:
 Q66HE1

 Cytogenetics:
 5q36

MW: 14.9 kDa

Gene Summary: Involved in nonsense-mediated mRNA decay (NMD) by acting as a bridge between the mRNA

decapping complex and the NMD machinery. May act by targeting the NMD machinery to the

P-body and recruiting the decapping machinery to aberrant mRNAs. Required for

UPF1/RENT1 localization to the P-body. Plays a role in glucocorticoid receptor-mediated mRNA degradation by interacting with the glucocorticoid receptor NR3C1 in a ligand-dependent manner when it is bound to the 5' UTR of target mRNAs and recruiting the RNA helicase UPF1 and the mRNA-decapping enzyme DCP1A, leading to RNA decay. Also acts as a

nuclear receptor coactivator. May play a role in controlling the energy balance between

energy storage and energy expenditure.[UniProtKB/Swiss-Prot Function]