

Product datasheet for RR205099L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hsd3b1 (NM_001007719) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hsd3b1 (NM_001007719) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Hsd3b1

Synonyms: MGC105549

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001007719

ORF Size: 1119 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR205099).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001007719.3</u>, <u>NP 001007720.3</u>

 RefSeq Size:
 1640 bp

 RefSeq ORF:
 1122 bp

 Locus ID:
 360348

 UniProt ID:
 P22071

 Cytogenetics:
 2q34





Gene Summary:

A bifunctional enzyme responsible for the oxidation and isomerization of 3beta-hydroxy-Delta(5)-steroid precursors to 3-oxo-Delta(4)-steroids, an essential step in steroid hormone biosynthesis. Specifically catalyzes the conversion of pregnenolone to progesterone, dehydroepiandrosterone (DHEA) to 4-androstenedione, and androstenediol to testosterone (PubMed:1537836, PubMed:1985917). Additionally, catalyzes the interconversion between 3beta-hydroxy and 3-oxo-5alpha-androstane steroids controlling the bioavalability of the active forms. Specifically converts dihydrotestosterone to its inactive form 5alpha-androstanediol, that does not bind androgen receptor/AR. Also converts androstanedione, a precursor of testosterone and estrone, to epiandrosterone (PubMed:1537836). Expected to use NAD(+) as preferred electron donor for the 3beta-hydroxy-steroid dehydrogenase activity and NADPH for the 3-ketosteroid reductase activity (Probable).[UniProtKB/Swiss-Prot Function]