

## Product datasheet for RR205806L3V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Large2 (NM\_199107) Rat Tagged ORF Clone Lentiviral Particle

### **Product data:**

Product Type: Lentiviral Particles

**Product Name:** Large2 (NM\_199107) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Large2
Synonyms: Gyltl1b

Mammalian Cell Puromycin

Selection:

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_199107

 ORF Size:
 2070 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RR205806).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** NM 199107.1, NP 954538.1

 RefSeq Size:
 2455 bp

 RefSeq ORF:
 2073 bp

 Locus ID:
 311202

 UniProt ID:
 Q6P7A1

 Cytogenetics:
 3q24







### **Gene Summary:**

Bifunctional glycosyltransferase with both xylosyltransferase and beta-1,3-glucuronyltransferase activities involved in the biosynthesis of the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan (DAG1). Phosphorylated O-mannosyl trisaccharid is required for binding laminin G-like domain-containing extracellular proteins with high affinity. Elongates the glucuronyl-beta-1,4-xylose-beta disaccharide primer structure by adding repeating units [-3-Xylose-alpha-1,3-GlcA-beta-1-] to produce a heteropolysaccharide. Has a higher activity toward alpha-dystroglycan than LARGE.[UniProtKB/Swiss-Prot Function]