

Product datasheet for RR205989L4

Auh (NM_001108407) Rat Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Auh (NM_001108407) Rat Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	Auh
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR205989).
Restriction Sites:	Sgfl-RsrII
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF.


ACCN:	NM_001108407
ORF Size:	759 bp



[View online »](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001108407.1 , NP_001101877.1
RefSeq Size:	1246 bp
RefSeq ORF:	762 bp
Locus ID:	361215
Cytogenetics:	17p14
Gene Summary:	Catalyzes the conversion of 3-methylglutaconyl-CoA to 3-hydroxy-3-methylglutaryl-CoA (By similarity). Also has itaconyl-CoA hydratase activity by converting itaconyl-CoA into citramalyl-CoA in the C5-dicarboxylate catabolism pathway (PubMed:13783048). The C5-dicarboxylate catabolism pathway is required to detoxify itaconate, a vitamin B12-poisoning metabolite (PubMed:13783048). Has very low enoyl-CoA hydratase activity (By similarity). Was originally identified as RNA-binding protein that binds in vitro to clustered 5'-AUUUA-3' motifs (By similarity).[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RR205989L4