

## Product datasheet for RR206980L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Syncrip (NM\_001047916) Rat Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Syncrip (NM\_001047916) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Syncrip

**Synonyms:** Ab2-339; hnRNP Q

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001047916

ORF Size: 1599 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RR206980).

Sequence:

Cytogenetics:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 001047916.1</u>, <u>NP 001041381.1</u>

8q31

 RefSeq Size:
 1648 bp

 RefSeq ORF:
 1602 bp

 Locus ID:
 363113

 UniProt ID:
 Q7TP47





## **Gene Summary:**

Heterogenous nuclear ribonucleoprotein (hnRNP) implicated in mRNA processing mechanisms. Component of the CRD-mediated complex that promotes MYC mRNA stability. Is associated in vitro with pre-mRNA, splicing intermediates and mature mRNA protein complexes. Binds to apoB mRNA AU-rich sequences. Part of the APOB mRNA editosome complex and may modulate the postranscriptional C to U RNA-editing of the APOB mRNA through either by binding to A1CF (APOBEC1 complementation factor), to APOBEC1 or to RNA itself. May be involved in translationally coupled mRNA turnover. Implicated with other RNA-binding proteins in the cytoplasmic deadenylation/translational and decay interplay of the FOS mRNA mediated by the major coding-region determinant of instability (mCRD) domain. Interacts in vitro preferentially with poly(A) and poly(U) RNA sequences. May be involved in cytoplasmic vesicle-based mRNA transport through interaction with synaptotagmins (By similarity).[UniProtKB/Swiss-Prot Function]