

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for RR207680L3V

Ncaph2 (NM_001024877) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Ncaph2 (NM_001024877) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Ncaph2
Synonyms:	MGC114417
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001024877
ORF Size:	1662 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR207680).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001024877.1, NP 001020048.1</u>
RefSeq Size:	2141 bp
RefSeq ORF:	1665 bp
Locus ID:	300149
UniProt ID:	<u>Q4V8I2</u>
Cytogenetics:	7q34



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Gene Summary:Regulatory subunit of the condensin-2 complex, a complex that seems to provide
chromosomes with an additional level of organization and rigidity and in establishing mitotic
chromosome architecture (By similarity). May promote the resolution of double-strand DNA
catenanes (intertwines) between sister chromatids. Condensin-mediated compaction likely
increases tension in catenated sister chromatids, providing directionality for type II
topoisomerase-mediated strand exchanges toward chromatid decatenation. Required for
decatenation of chromatin bridges at anaphase. Early in neurogenesis, may play an essential
role to ensure accurate mitotic chromosome condensation in neuron stem cells, ultimately
affecting neuron pool and cortex size (By similarity). Seems to have lineage-specific role in T-
cell development (By similarity).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US