

# Product datasheet for RR208578L4V

## Adrm1 (NM\_031708) Rat Tagged ORF Clone Lentiviral Particle

### **Product data:**

#### **Product Type: Lentiviral Particles Product Name:** Adrm1 (NM\_031708) Rat Tagged ORF Clone Lentiviral Particle Symbol: Adrm1 Gp110 Synonyms: **Mammalian Cell** Puromycin Selection: Vector: pLenti-C-mGFP-P2A-Puro (PS100093) mGFP Tag: NM 031708 ACCN: ORF Size: 1221 bp The ORF insert of this clone is exactly the same as(RR208578). **ORF** Nucleotide Sequence: **OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info **OTI** Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. **RefSeq:** NM 031708.1, NP 113896.1 **RefSeq Size:** 1444 bp **RefSeq ORF:** 1224 bp Locus ID: 65138 **UniProt ID:** Q9JMB5 Cytogenetics: 3q43



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



Gene Summary: Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. Within the complex, functions as a proteasomal ubiquitin receptor. Engages and thus activates 19S-associated deubiquitinases UCHL5 and PSMD14 during protein degradation. UCHL5 reversibly associate with the 19S regulatory particle whereas PSMD14 is an intrinsic subunit of the proteasome lid subcomplex.[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US