

Product datasheet for RR208581L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Hbs1l (NM_001011934) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Hbs1l (NM_001011934) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Hbs1l

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM_001011934

ORF Size: 2037 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RR208581).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 001011934.1, NP 001011934.1

 RefSeq Size:
 2721 bp

 RefSeq ORF:
 2040 bp

 Locus ID:
 293408

 UniProt ID:
 Q6AXM7

Cytogenetics: 1p12







Gene Summary:

Cotranslational quality control factor involved in the No-Go Decay (NGD) pathway. In the presence of ABCE1 and PELO, is required for 48S complex formation from 80S ribosomes and dissociation of vacant 80S ribosomes. Together with PELO and in presence of ABCE1, recognizes stalled ribosomes and promotes dissociation of elongation complexes assembled on non-stop mRNAs; this triggers endonucleolytic cleavage of the mRNA, a mechanism to release non-functional ribosomes and to degrade damaged mRNAs as part of the No-Go Decay (NGD) pathway.[UniProtKB/Swiss-Prot Function]