

Product datasheet for RR209465

Ppargc1b (NM_176075) Rat Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ppargc1b (NM_176075) Rat Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Ppargc1b
Synonyms:	Perc; PGC1beta
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF


ACCN:	NM_176075
ORF Size:	2970 bp



[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_176075.2 , NP_788264.1
RefSeq Size:	3163 bp
RefSeq ORF:	2973 bp
Locus ID:	291567
UniProt ID:	Q811R2
Cytogenetics:	18q12.1
MW:	109.4 kDa
Gene Summary:	Plays a role of stimulator of transcription factors and nuclear receptors activities. Activates transcriptional activity of estrogen receptor alpha, nuclear respiratory factor 1 (NRF1) and glucocorticoid receptor in the presence of glucocorticoids. May play a role in constitutive non-adrenergic-mediated mitochondrial biogenesis as suggested by increased basal oxygen consumption and mitochondrial number when overexpressed. May be part of the pathways regulating the elevation of gluconeogenesis, beta-oxidation of fatty acids and ketogenesis during fasting. Stimulates SREBP-mediated lipogenic gene expression in the liver. Induces energy expenditure and antagonizes obesity when overexpressed. Induces also the expression of mitochondrial genes involved in oxidative metabolism. Induces the expression of PERM1 in the skeletal muscle in an ESRRA-dependent manner.[UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RR209465