


Product datasheet for **RR209469L3**

Fbxw5 (NM_001025730) Rat Tagged Lenti ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Fbxw5 (NM_001025730) Rat Tagged Lenti ORF Clone
Tag:	Myc-DDK
Symbol:	Fbxw5
Synonyms:	MGC116348
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR209469).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:




* The last codon before the Stop codon of the ORF.

ACCN:	NM_001025730
ORF Size:	1707 bp



OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001025730.2 , NP_001020901.1
RefSeq Size:	2345 bp
RefSeq ORF:	1710 bp
Locus ID:	362081
UniProt ID:	Q4KLI9
Cytogenetics:	3p13
Gene Summary:	Substrate recognition component of both SCF (SKP1-CUL1-F-box protein) and DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complexes. Substrate recognition component of the SCF(FBXW5) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of SASS6 during S phase, leading to prevent centriole reduplication. The SCF(FBXW5) complex also mediates ubiquitination and degradation of actin-regulator EPS8 during G2 phase, leading to the transient degradation of EPS8 and subsequent cell shape changes required to allow mitotic progression. Substrate-specific adapter of the DCX(FBXW5) E3 ubiquitin-protein ligase complex which mediates the polyubiquitination and subsequent degradation of TSC2. May also act as a negative regulator of MAP3K7/TAK1 signaling in the interleukin-1B (IL1B) signaling pathway (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RR209469L3