

Product datasheet for RR209469L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Fbxw5 (NM_001025730) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Fbxw5 (NM 001025730) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Fbxw5

Synonyms: MGC116348

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001025730

ORF Size: 1707 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR209469).

Sequence:

UniProt ID:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001025730.2</u>, <u>NP 001020901.1</u>

Q4KLI9

 RefSeq Size:
 2345 bp

 RefSeq ORF:
 1710 bp

 Locus ID:
 362081

Cytogenetics: 3p13





Gene Summary:

Substrate recognition component of both SCF (SKP1-CUL1-F-box protein) and DCX (DDB1-CUL4-X-box) E3 ubiquitin-protein ligase complexes. Substrate recognition component of the SCF(FBXW5) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of SASS6 during S phase, leading to prevent centriole reduplication. The SCF(FBXW5) complex also mediates ubiquitination and degradation of actin-regulator EPS8 during G2 phase, leading to the transient degradation of EPS8 and subsequent cell shape changes required to allow mitotic progression. Substrate-specific adapter of the DCX(FBXW5) E3 ubiquitin-protein ligase complex which mediates the polyubiquitination and subsequent degradation of TSC2. May also act as a negative regulator of MAP3K7/TAK1 signaling in the interleukin-1B (IL1B) signaling pathway (By similarity). [UniProtKB/Swiss-Prot Function]