

Product datasheet for RR209542

Jmjd6 (NM_001012143) Rat Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Jmjd6 (NM_001012143) Rat Tagged ORF Clone

Tag: Myc-DDK

Symbol: Jmjd6

Synonyms: Ptdsr

Mammalian Cell

Neomycin

Selection: Vector:

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

ORF Nucleotide >RR209542 representing NM_001012143
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGAACCACAAGAGCAAGAAGCGCATCCGCGAGGCCAAGCGGAGTGCGCGGCCCGAGCTCAAGGACTCGC TTGACTGGACGCGACACAATTACTATGAGAGCTACCCGCTGAGCCCTGCGGCCGTGCCGGATAACGTGGA GAGAGCTGATGCTTTGCAGCTGTCGGTGAAAGAATTTGTAGAGCGGTATGAAAGACCTTACAAGCCCGTG GTTCTGCTGAATGCACAAGAGGGCTGGTCCGCACAGGAGAAATGGACTCTGGAGCGTCTCAAAAGGAAAT ACCGGAACCAGAAGTTCAAATGCGGTGAGGATAATGACGGCTACTCGGTGAAGATGAAAATGAAATATTA CATCGAGTACATGGAGAGCACCCGCGATGACAGTCCCCTTTACATCTTTGATAGCAGCTATGGTGAACAC CCCAAAAGAAGGAAACTTTTGGAAGACTATAAGGTGCCCAAGTTTTTCACAGATGATCTTTTCCAATATG CGGGGGAGAAGCGCAGACCCCCTTACAGGTGGTTTGTGATGGGGCCACCACGTTCTGGAACTGGGATTCA CATTGACCCTCTGGGGACCAGTGCCTGGAATGCCTTAGTTCAGGGTCACAAGCGGTGGTGCCTGTTCCCA ACAAACACCCCGAGAACTCATCAAGGTGACCCGAGAAGAAGGAGGAAACCAACAGGATGAAGCAATTA CCTGGTTTAATGTCATTTATCCCCGGACACAGCTTCCAACCTGGCCACCCGAATTCAAACCCCTGGAGAT ATTGCAGAAACCAGGAGAAACTGTCTTTGTACCAGGGGGCTGGTGGCACGTCGTCCTCAATCTTGACACC ACCATTGCCATCACCCAGAACTTTGCCAGCAGCACCAACTTCCCTGTTGTGTGGCACAAGACGGTAAGAG GGCGGCCAAAGTTATCCAGGAAGTGGTATAGGATCTTGAAGCAGGAGCACCCTGAGCTGGCAGTGCTTGC CGACGCTGTTGACCTCCAAGAGTCCACAGGCATCGCCTCTGACAGCTCCAGTGACTCTTCTAGCTCCTCC AGCTCCAGCTCGGACTCAGACTCGGAGTGTGAATCTGGGTCAGAAGGTGATGGGACAACACACCACA GGAAGAAGAGGAGAACTTGCAGCATGGTGGGAAATGGGGACACTACCTCACAGGATGACTGCGTGAGCAA AGAGCGCAGTTCCTCCAGG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Protein Sequence: >RR209542 representing NM_001012143

Red=Cloning site Green=Tags(s)

MNHKSKKRIREAKRSARPELKDSLDWTRHNYYESYPLSPAAVPDNVERADALQLSVKEFVERYERPYKPV VLLNAQEGWSAQEKWTLERLKRKYRNQKFKCGEDNDGYSVKMKMKYYIEYMESTRDDSPLYIFDSSYGEH PKRRKLLEDYKVPKFFTDDLFQYAGEKRRPPYRWFVMGPPRSGTGIHIDPLGTSAWNALVQGHKRWCLFP TNTPRELIKVTREEGGNQQDEAITWFNVIYPRTQLPTWPPEFKPLEILQKPGETVFVPGGWWHVVLNLDT TIAITQNFASSTNFPVVWHKTVRGRPKLSRKWYRILKQEHPELAVLADAVDLQESTGIASDSSSDSSSSS SSSSSDSDSECESGSEGDGTTHRRKKRRTCSMVGNGDTTSQDDCVSKERSSSR


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites:

Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the OR

ACCN: NM_001012143

ORF Size: 1209 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeg: NM 001012143.2, NP 001012143.2

 RefSeq Size:
 1730 bp

 RefSeq ORF:
 1212 bp

 Locus ID:
 360665

 UniProt ID:
 Q6AYK2

 Cytogenetics:
 10q32.2


 MW:
 46.5 kDa

Gene Summary:

Dioxygenase that can both act as a arginine demethylase and a lysyl-hydroxylase. Acts as a lysyl-hydroxylase that catalyzes 5-hydroxylation on specific lysine residues of target proteins such as U2AF2/U2AF65 and LUC7L2. Regulates RNA splicing by mediating 5-hydroxylation of U2AF2/U2AF65, affecting the pre-mRNA splicing activity of U2AF2/U2AF65. Hydroxylates its own N-terminus, which is required for homooligomerization. In addition to peptidyl-lysine 5dioxygenase activity, may act as an RNA hydroxylase, as suggested by its ability to bind single strand RNA. Also acts as an arginine demethylase which preferentially demethylates asymmetric dimethylation. Demethylates histone H3 at 'Arg-2' (H3R2me) and histone H4 at 'Arg-3' (H4R3me), including mono-, symmetric di- and asymmetric dimethylated forms, thereby playing a role in histone code. However, histone arginine demethylation may not constitute the primary activity in vivo. In collaboration with BRD4, interacts with the positive transcription elongation factor b (P-TEFb) complex in its active form to regulate polymerase II promoter-proximal pause release for transcriptional activation of a large cohort of genes. On distal enhancers, so called anti-pause enhancers, demethylates both histone H4R3me2 and the methyl cap of 7SKsnRNA leading to the dismissal of the 7SKsnRNA:HEXIM1 inhibitor complex. After removal of repressive marks, the complex BRD4:JMJD6 attract and retain the P-TEFb complex on chromatin, leading to its activation, promoter-proximal polymerase II pause release, and transcriptional activation. Demethylates other arginine methylatedproteins such as ESR1. Has no histone lysine demethylase activity (By similarity). Required for differentiation of multiple organs during embryogenesis. Acts as a key regulator of hematopoietic differentiation: required for angiogenic sprouting by regulating the pre-mRNA splicing activity of U2AF2/U2AF65 (By similarity). Seems to be necessary for the regulation of macrophage cytokine responses (By similarity).[UniProtKB/Swiss-Prot Function]



Product images:



Circular map for RR209542