

Product datasheet for **RR212516L4V**

Dhx36 (NM_001107678) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	Dhx36 (NM_001107678) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Dhx36
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001107678
ORF Size:	3000 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR212516).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_001107678.1 , NP_001101148.1
RefSeq Size:	5664 bp
RefSeq ORF:	3003 bp
Locus ID:	310461
UniProt ID:	D4A2Z8
Cytogenetics:	2q31



[View online »](#)

Gene Summary:

Multifunctional ATP-dependent helicase that unwinds G-quadruplex (G4) structures (By similarity). Plays a role in many biological processes such as genomic integrity, gene expression regulations and as a sensor to initiate antiviral responses (PubMed:23651854). G4 structures correspond to helical structures containing guanine tetrads (By similarity). Binds with high affinity to and unwinds G4 structures that are formed in nucleic acids (G4-ADN and G4-RNA) (By similarity). Plays a role in genomic integrity. Converts the G4-RNA structure present in telomerase RNA template component (TREC) into a double-stranded RNA to promote P1 helix formation that acts as a template boundary ensuring accurate reverse transcription (By similarity). Plays a role in transcriptional regulation. Resolves G4-DNA structures in promoters of genes, such as YY1, KIT/c-kit and ALPL and positively regulates their expression (By similarity). Plays a role in post-transcriptional regulation. Unwinds a G4-RNA structure located in the 3' UTR polyadenylation site of the pre-mRNA TP53 and stimulates TP53 pre-mRNA 3'-end processing in response to ultraviolet (UV)-induced DNA damage (By similarity). Binds to the precursor-microRNA-134 (pre-miR-134) terminal loop and regulates its transport into the synapto-dendritic compartment (PubMed:23651854). Involved in the pre-miR-134-dependent inhibition of target gene expression and the control of dendritic spine size (PubMed:23651854). Plays a role in the regulation of cytoplasmic mRNA translation and mRNA stability. Binds to both G4-RNA structures and alternative non-quadruplex-forming sequence within the 3' UTR of the PITX1 mRNA regulating negatively PITX1 protein expression. Binds to both G4-RNA structure in the 5'-UTR and AU-rich elements (AREs) localized in the 3' UTR of NKX2-5 mRNA to either stimulate protein translation or induce mRNA decay in an ELAVL1-dependent manner, respectively. Binds also to ARE sequences present in several mRNAs mediating exosome-mediated 3'-5' mRNA degradation. Involved in cytoplasmic urokinase-type plasminogen activator (uPA) mRNA decay (By similarity). Component of a multi-helicase-TICAM1 complex that acts as a cytoplasmic sensor of viral double-stranded RNA (dsRNA) and plays a role in the activation of a cascade of antiviral responses including the induction of proinflammatory cytokines via the adapter molecule TICAM1. Required for the early embryonic development and hematopoiesis. Involved in the regulation of cardioblast differentiation and proliferation during heart development. Involved in spermatogonia differentiation. May play a role in ossification (By similarity).[UniProtKB/Swiss-Prot Function]