

## Product datasheet for **RR213421L4V**

### Ficd (NM\_001010946) Rat Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Ficd (NM_001010946) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Ficd
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
Tag:	mGFP
ACCN:	NM_001010946
ORF Size:	1374 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR213421).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001010946.1</a> , <a href="#">NP_001010946.1</a>
RefSeq Size:	1965 bp
RefSeq ORF:	1377 bp
Locus ID:	288741
UniProt ID:	<a href="#">Q6AY47</a>
Cytogenetics:	12q16



[View online »](#)

**Gene Summary:**

Protein that can both mediate the addition of adenosine 5'-monophosphate (AMP) to specific residues of target proteins (AMPylation), and the removal of the same modification from target proteins (de-AMPylation), depending on the context (By similarity). The side chain of Glu-231 determines which of the two opposing activities (AMPylase or de-AMPylase) will take place (By similarity). Acts as a key regulator of the ERN1/IRE1-mediated unfolded protein response (UPR) by mediating AMPylation or de-AMPylation of HSPA5/BiP (By similarity). In unstressed cells, acts as an adenylyltransferase by mediating AMPylation of HSPA5/BiP at 'Thr-518', thereby inactivating it (By similarity). In response to endoplasmic reticulum stress, acts as a phosphodiesterase by mediating removal of ATP (de-AMPylation) from HSPA5/BiP at 'Thr-518', leading to restore HSPA5/BiP activity (By similarity). Although it is able to AMPylate RhoA, Rac and Cdc42 Rho GTPases in vitro, Rho GTPases do not constitute physiological substrates (By similarity).[UniProtKB/Swiss-Prot Function]