

## Product datasheet for RR213833L3V

**OriGene Technologies, Inc.**  
 9620 Medical Center Drive, Ste 200  
 Rockville, MD 20850, US  
 Phone: +1-888-267-4436  
<https://www.origene.com>  
[techsupport@origene.com](mailto:techsupport@origene.com)  
 EU: [info-de@origene.com](mailto:info-de@origene.com)  
 CN: [techsupport@origene.cn](mailto:techsupport@origene.cn)

### Dis3l (NM\_001008380) Rat Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Dis3l (NM_001008380) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Dis3l
Synonyms:	RGD1308959
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001008380
ORF Size:	2913 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR213833).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_001008380.1</a> , <a href="#">NP_001008381.1</a>
RefSeq Size:	3234 bp
RefSeq ORF:	2916 bp
Locus ID:	363077
UniProt ID:	<a href="#">Q5U2P0</a>
Cytogenetics:	8q24



**Gene Summary:**

Putative cytoplasm-specific catalytic component of the RNA exosome complex which has 3'->5' exoribonuclease activity and participates in a multitude of cellular RNA processing and degradation events. In the cytoplasm, the RNA exosome complex is involved in general mRNA turnover and specifically degrades inherently unstable mRNAs containing AU-rich elements (AREs) within their 3' untranslated regions, and in RNA surveillance pathways, preventing translation of aberrant mRNAs. It seems to be involved in degradation of histone mRNA. [UniProtKB/Swiss-Prot Function]