

Product datasheet for RR213834L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nfatc4 (NM_001107264) Rat Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Nfatc4 (NM 001107264) Rat Tagged ORF Clone Lentiviral Particle

Symbol: Nfatc4

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001107264

ORF Size: 2703 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RR213834).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001107264.1</u>, <u>NP 001100734.1</u>

 RefSeq Size:
 3170 bp

 RefSeq ORF:
 2706 bp

 Locus ID:
 305897

 UniProt ID:
 D3Z9H7

 Cytogenetics:
 15p13





Gene Summary:

Ca(2+)-regulated transcription factor that is involved in several processes, including the development and function of the immune, cardiovascular, musculoskeletal, and nervous systems. Involved in T-cell activation, stimulating the transcription of cytokine genes, including that of IL2 and IL4 (PubMed:12370307). Along with NFATC3, involved in embryonic heart development. Involved in mitochondrial energy metabolism required for cardiac morphogenesis and function. Transactivates many genes involved in heart physiology. Along with GATA4, binds to and activates NPPB/BNP promoter (PubMed:9568714). Activates NPPA/ANP/ANF and MYH7/beta-MHC transcription (PubMed:19026640). Binds to and transactivates AGTR2 gene promoter. Involved in the regulation of adult hippocampal neurogenesis. Involved in BDNF-driven pro-survival signaling in hippocampal adult-born neurons. Involved in the formation of long-term spatial memory and long-term potentiation. In cochlear nucleus neurons, may play a role in deafferentation-induced apoptosis during a developmental critical period when auditory neurons depend on afferent input for survival (By similarity). Binds to and activates the BACE1/Beta-secretase 1 promoter, hence may regulate the proteolytic processing of the amyloid precursor protein (APP). Plays a role in adipocyte differentiation. May be involved in myoblast differentiation into myotubes (By similarity). Binds the consensus DNA sequence 5'-GGAAAAT-3' (By similarity). In the presence of CREBBP, activates TNF transcription. Binds to PPARG gene promoter and regulates its activity (By similarity). Binds to PPARG and REG3G gene promoters (By similarity). [UniProtKB/Swiss-Prot Function]