

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RR214224L3V

## Lrrc8c (NM\_001037179) Rat Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	Lrrc8c (NM_001037179) Rat Tagged ORF Clone Lentiviral Particle
Symbol:	Lrrc8c
Synonyms:	RGD1306585
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_001037179
ORF Size:	2409 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RR214224).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 001037179.1, NP 001032256.1</u>
RefSeq Size:	2883 bp
RefSeq ORF:	2412 bp
Locus ID:	289443
UniProt ID:	<u>Q498T9</u>
Cytogenetics:	14p22



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Lrrc8c (NM_001037179) Rat Tagged ORF Clone Lentiviral Particle – RR214224L3V
Gene Summary:	Non-essential component of the volume-regulated anion channel (VRAC, also named VSOAC
	channel), an anion channel required to maintain a constant cell volume in response to

channel), an anion channel required to maintain a constant cell volume in response to extracellular or intracellular osmotic changes (PubMed:28833202). The VRAC channel conducts iodide better than chloride and can also conduct organic osmolytes like taurine (By similarity). Plays a redundant role in the efflux of amino acids, such as aspartate and glutamate, in response to osmotic stress. Channel activity requires LRRC8A plus at least one other family member (LRRC8B, LRRC8C, LRRC8D or LRRC8E); channel characteristics depend on the precise subunit composition (PubMed:28833202).[UniProtKB/Swiss-Prot Function]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US