

Product datasheet for **RR214781**

Rnf146 (NM_001012060) Rat Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: Rnf146 (NM_001012060) Rat Tagged ORF Clone
Tag: Myc-DDK
Symbol: Rnf146
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
ORF Nucleotide Sequence: >RR214781 representing NM_001012060
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGAATGATGGCCGGCTGTGGTGAATATGATCACTCACTAAATATGCTTCTACCAATAAGAAGCGCA
GTGAGACCTGTTCTAACACTGCACCTTCTCTAACAGTTCCCGAGTGTGCCATTTGTCTACAACATGTGT
TCATCCAGTCAGTCTGCCCTGTAAGCATGTTTTCTGTTATCTGTGTGTAAGGGCGCTTCATGGCTCGGG
AAGCGATGTGCTCTTTGTCGGCAAGAGATTCCTGAGGATTTCTTGACAAGCCAACCTTGTGTACCAG
AAGAACTTAAGGCTGCAAGCAGAGGAAATGGTGAATATGTGTGGTATTATGAAGGAAGAAATGGATGGTG
GCAGTATGATGAGCGCACCAGTCGGGAGCTAGAAGATGCTTTTTCCAAAGGTAAAAAGAACCGAAATG
TTAATTGCTGGATTTCTGTACGTTGCTGATCTTGAACATGGTTCAATATAGGAGAAATGAACATGGAC
GTCGCAGAAAGATTAAGAGATATAATAGATATACCAAAGAAGGGAGTGGCTGGACTTCGGCTGGACTG
TGACAGCAACTGTAATCTAGCCAGAGAGATTCTGCCGATGGTGGGACAGTGGTCCAGCACACACT
GGAGCTTCTGTGAGCTTCCAGTGCCATCTTCTACAAGGCCTCTAACATCAGTCGATGGTCAGTTAACCA
GCCCTGTAACACCGTCCCCTGATGCAGGGCCTTTTGAAGACTCTTTGCTCATTACAACCTCAGTGG
AGACAGCATAGCTGAAAGGAGTCATAGAGGTGAAGGAGAAGAAGTACCGAATCACCATCTTCTGGTAGA
GTACCAGATACGTCCACTGAAGAAACAGAGTCGGATGCCAGTAGTGATATTGAGGATGCCCTGTGGTGG
TTGCACAGCACTCCTTGACCCAACAGAGACTTTTGGTTTCAAGTGCAAATCAGACAGTAGCTGAACGGTC
TGACCGACCAGTTGCAGGGGGTGGGACCATGAGTGTCAATGTCAGATCCAGAAGGCCTGACGGACAGTGC
ACAGTGACAGAGGTT

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)


Protein Sequence: >RR214781 representing NM_001012060
 Red=Cloning site Green=Tags(s)

MEMMAGCGEIDHSLNMLPTNKKASETCSNTAPSLTVPECAICLQTCVHPVSLPCKHVFCYLCVKGASWLG
 KRCALCRQEIPEDFLDKPTLLSPEELKAASRNGEYVWYVEGRNGWQYDERTSRELEDAFSKGGKNTM
 LIAGFLYVADLENMVQYRRNEHGRRRKIKRDIIDIPKKGAVGLRLDCDSNTVNLARESSADGADSGSAHT
 GASVQLPVPSSSTRPLTSVDGQLTSPVTPSPDAGASLEDSFAHLQLSGDSTIAERSHRGEEDHESPSSGR
 VPDTSTEETESDASSDIEDAPVVVAQHSLTQQRLLVSSANQTVAESDRPVAGGGTMSVNVRSRRPDGQC
 TVTEV


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI

Cloning Scheme:



Plasmid Map:



ACCN: NM_001012060

ORF Size:	1065 bp
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001012060.2 , NP_001012060.2
RefSeq Size:	1889 bp
RefSeq ORF:	1068 bp
Locus ID:	308051
UniProt ID:	Q5XIK5
Cytogenetics:	1p11
MW:	38.6 kDa

Gene Summary:

E3 ubiquitin-protein ligase that specifically binds poly-ADP-ribosylated (PARsylated) proteins and mediates their ubiquitination and subsequent degradation. May regulate many important biological processes, such as cell survival and DNA damage response. Acts as an activator of the Wnt signaling pathway by mediating the ubiquitination of PARsylated AXIN1 and AXIN2, 2 key components of the beta-catenin destruction complex. Acts in cooperation with tankyrase proteins (TNKS and TNKS2), which mediate PARsylation of target proteins AXIN1, AXIN2, BLZF1, CASC3, TNKS and TNKS2. Recognizes and binds tankyrase-dependent PARsylated proteins via its WWE domain and mediates their ubiquitination. May regulate TNKS and TNKS2 subcellular location, preventing aggregation at a centrosomal location. Neuroprotective protein. Protects the brain against N-methyl-D-aspartate (NMDA) receptor-mediated glutamate excitotoxicity and ischemia, by interfering with PAR-induced cell death, called parthanatos. Prevents nuclear translocation of AIFM1 in a PAR-binding dependent manner. Does not affect PARP1 activation. Protects against cell death induced by DNA damaging agents, such as N-methyl-N-nitro-N-nitrosoguanidine (MNNG) and rescues cells from G1 arrest. Promotes cell survival after gamma-irradiation. Facilitates DNA repair. Neuroprotective protein. Protects the brain against N-methyl-D-aspartate (NMDA) receptor-mediated glutamate excitotoxicity and ischemia, by interfering with PAR-induced cell death, called parthanatos. Prevents nuclear translocation of AIFM1 in a PAR-binding dependent manner. Does not affect PARP1 activation (By similarity).[UniProtKB/Swiss-Prot Function]