

## Product datasheet for **RR215712**

### Oaz2 (NM\_001301297) Rat Tagged ORF Clone

#### Product data:

Product Type: Expression Plasmids  
Product Name: Oaz2 (NM\_001301297) Rat Tagged ORF Clone  
Tag: Myc-DDK  
Symbol: Oaz2  
Synonyms: RGD1562933  
Vector: pCMV6-Entry (PS100001)  
E. coli Selection: Kanamycin (25 ug/mL)  
Cell Selection: Neomycin  
ORF Nucleotide Sequence: >RR215712 representing NM\_001301297  
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGC**

ATGATAAACACCCAGGACAGTATTTGCCTTTGAGTAAGTGTCCCGAGCTCCAGTGCTGCAGGCACATTG  
TTCCAGGGCCTCTGTGGTGCTCCATGATAAACACCCAGGACAGTATTTGCCTTTGAGTAAGTGTCCCA  
GCTCCAGTGCTGCAGGCACATTGTTCCAGGGCCTCTGTGGTGCTCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RR215712 representing NM\_001301297  
Red=Cloning site Green=Tags(s)

MINTQDSILPLSNCPQLQCCRHIIVGPLWCSMINTQDSILPLSNCPQLQCCRHIIVGPLWCS


**TR**TRPLE**QKL**ISEEDLAANDILDYKDDDDKV

Restriction Sites: SgfI-MluI




[View online »](#)

## Cloning Scheme:



## Plasmid Map:



ACCN: NM\_001301297

ORF Size: 183 bp

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">NM_001301297.1</a> , <a href="#">NP_001288226.1</a>
<b>RefSeq Size:</b>	1825 bp
<b>RefSeq ORF:</b>	568 bp
<b>Locus ID:</b>	501454
<b>Cytogenetics:</b>	8q24
<b>MW:</b>	6.9 kDa
<b>Gene Summary:</b>	<p>The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family, which plays a role in cell growth and proliferation by regulating intracellular polyamine levels. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 2, the second member of the antizyme family. Like antizyme 1, antizyme 2 has broad tissue distribution, inhibits ODC activity and polyamine uptake, and stimulates ODC degradation in vivo; however, it fails to promote ODC degradation in vitro. Antizyme 2 is expressed at lower levels than antizyme 1, but is evolutionary more conserved, suggesting it likely has an important biological role. Studies also show different subcellular localization of antizymes 1 and 2, indicating specific function for each antizyme in discrete compartments of the cell. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2014]</p>