

Product datasheet for RR215898

Cebpb (NM 001301720) Rat Tagged ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Cebpb (NM_001301720) Rat Tagged ORF Clone

Tag: Myc-DDK
Symbol: Cebpb

Synonyms: Il6dbp; NF-IL6; TCF5

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RR215898 representing NM_001301720
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RR215898 representing NM_001301720

Red=Cloning site Green=Tags(s)

MAAGFPFALRAYLGYQATPSGSSGSLSTSSSSSPPGTPSPADAKAAPAACFAGPPAAPAKAKAKKAVDKL SDEYKMRRERNNIAVRKSRDKAKMRNLETQHKVLELTAENERLQKKVEQLSRELSTLRNLFKQLPEPLLA

SAGHC

TRTRPLEQKLISEEDLAANDILDYKDDDDK**V**

Restriction Sites: Sgfl-Mlul



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

Plasmid Map:



ACCN: NM_001301720

ORF Size: 435 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001301720.1</u>, <u>NP 001288649.1</u>

 RefSeq Size:
 1475 bp

 RefSeq ORF:
 438 bp

 Locus ID:
 24253

 UniProt ID:
 P21272

 Cytogenetics:
 3q42

 MW:
 15.6 kDa

Gene Summary: This intronless gene encodes a member of the transcription factor family whose members

contain a basic leucine-zipper domain. The encoded protein functions as a homodimer but can also form heterodimers with CCAAT/enhancer-binding proteins alpha, delta, and gamma.

The encoded protein plays important roles in several cellular processes and in various

diseases, including regulating cell proliferation, differentiation, apoptosis and

neuroinflammation, and being involved in brain injury and inflammatory progression. The use of alternative in-frame AUG start codons results in multiple protein isoforms, each with different cellular localizations and distinct biological functions. [provided by RefSeq, Sep 2014]