

# **Product datasheet for RR215899**

## Vegfa (NM\_001287110) Rat Tagged ORF Clone

#### **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Vegfa (NM\_001287110) Rat Tagged ORF Clone

Tag: Myc-DDK
Symbol: Vegfa

**Synonyms:** Vegf; VEGF-A; VEGF111; VEGF164; VPF

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

ORF Nucleotide >RR215899 representing NM\_001287110
Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGAGATCTGCC

GCCGCGATCGCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RR215899 representing NM\_001287110

Red=Cloning site Green=Tags(s)

MNFLLSWVHWTLALLLYLHHAKWSQAAPTTEGEQKAHEVVKFMDVYQRSYCRPIETLVDIFQEYPDEIEY IFKPSCVPLMRCAGCCNDEALECVPTSESNVTMQIMRIKPHQSQHIGEMSFLQHSRCECRPKKDRTKPEK

**CDKPRR** 

**TRTRPL**EQKLISEEDLAANDILDYKDDDDK**V** 

**Restriction Sites:** Sgfl-Mlul




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




#### **Cloning Scheme:**





<sup>\*</sup> The last codon before the Stop codon of the ORF

### Plasmid Map:



**ACCN:** NM\_001287110

ORF Size: 438 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001287110.1</u>, <u>NP 001274039.1</u>

 RefSeq Size:
 3342 bp

 RefSeq ORF:
 441 bp

 Locus ID:
 83785

 UniProt ID:
 P16612

 Cytogenetics:
 9q12

 MW:
 17.2 kDa

**Gene Summary:** This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding

protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is

isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. [provided by RefSeq, Nov 2015]

produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some