


Product datasheet for RR216521

Slc30a7 (NM_001191715) Rat Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
 Product Name: Slc30a7 (NM_001191715) Rat Tagged ORF Clone
 Tag: Myc-DDK
 Symbol: Slc30a7
 Mammalian Cell Selection: Neomycin
 Vector: pCMV6-Entry (PS100001)
 E. coli Selection: Kanamycin (25 ug/mL)
 Restriction Sites: SgfI-MluI
 Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF


ACCN: NM_001191715
 ORF Size: 1134 bp



[View online >](#)

OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001191715.1 , NP_001178644.1
RefSeq Size:	2032 bp
RefSeq ORF:	1137 bp
Locus ID:	310801
UniProt ID:	Q5BJM8
Cytogenetics:	2q41
MW:	41.8 kDa
Gene Summary:	Seems to facilitate zinc transport from the cytoplasm into the Golgi apparatus. Partly regulates cellular zinc homeostasis. Required with ZNT5 for the activation of zinc-requiring enzymes, alkaline phosphatases (ALPs). Transports zinc into the lumens of the Golgi apparatus and the vesicular compartments where ALPs locate, thus, converting apoALPs to holoALPs. Required with ZNT5 and ZNT6 for the activation of TNAP (By similarity). [UniProtKB/Swiss-Prot Function]

Product images:



Circular map for RR216521