

## Product datasheet for SA6035

### CD158i / KIR2DS4 Human Protein

#### Product data:

|                                       |   |
|---------------------------------------|---|
| Product Type:                         | Recombinant Proteins  |
| Description:                          | CD158i / KIR2DS4 human recombinant protein, 0.1 mg  |
| Species:                              | Human   |
| Expression Host:                      | E. coli   |
| Expression cDNA Clone or AA Sequence: | MEGVHRKPSF LALPGHLVKS EETVILQCWS DVMFEHLLH REGKFNNTLH LIGEHHDGVS<br>KANFSIGPMM PVLAGTYRCY GSVPHSPYQL SAPSDPLDMV IIGLYEKPSL SAQPGPTVQA<br>GENVTLS CSS RSSYDMYHLS REGEAHERRL PAVRSINGTF QADFPLGPAT HGGTYRCFGS<br>FRDAPYEWSN SSDPLLVSVT GN |
| Predicted MW:                         | 22.2 kDa  |
| Concentration:                        | lot specific  |
| Purity:                               | >95% by SDS-PAGE  |
| Buffer:                               | Presentation State: Purified<br>State: Liquid protein<br>Buffer System: 20 mM Tris-HCl buffer (pH 7.5)  |
| Preparation:                          | Liquid protein  |
| Protein Description:                  | The recombinant KIR2DS4 protein was purified by FPLC gel-filtration chromatography, after refolding of the isolated inclusion bodies in a redox buffer.   |
| Storage:                              | Store at 2 - 8 °C for up to one month or (in aliquots) at -20 °C. Avoid repeated freezing and thawing.  |
| Stability:                            | Shelf life: one year from despatch.   |
| RefSeq:                               | <a href="#">NP_001268900</a>  |
| Locus ID:                             | 3809  |
| UniProt ID:                           | <a href="#">P43632</a> , <a href="#">A0A0C4ZN30</a> , <a href="#">Q8NHJ0</a>  |
| Cytogenetics:                         | 19q13.42  |
| Synonyms:                             | CD158I; KIR-2DS4; KIR1D; KIR412; KKA3; NKAT-8; NKAT8  |



[View online »](#)

**Summary:**

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]

**Protein Families:**

Transmembrane

**Protein Pathways:**

Antigen processing and presentation, Natural killer cell mediated cytotoxicity

**Product images:**