

Product datasheet for SA6035X

CD158i / KIR2DS4 Human Protein

Product data:

Product Type:	Recombinant Proteins
Description:	CD158i / KIR2DS4 human protein, 0.5 mg
Species:	Human
Expression Host:	E. coli
Predicted MW:	22.2 kDa
Purity:	>95%
Buffer:	Presentation State: Purified
RefSeq:	NP_001268900
Locus ID:	3809
UniProt ID:	P43632 , A0A0C4ZN30 , Q8NHJ0
Cytogenetics:	19q13.42
Synonyms:	CD158i; KIR-2DS4; KIR1D; KIR412; KKA3; NKAT-8; NKAT8
Summary:	Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]
Protein Families:	Transmembrane
Protein Pathways:	Antigen processing and presentation, Natural killer cell mediated cytotoxicity



[View online »](#)

Product images: