

## Product datasheet for **SC101215**

### **KRTCAP2 (NM\_173852) Human Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	KRTCAP2 (NM_173852) Human Untagged Clone
Tag:	Tag Free
Symbol:	KRTCAP2
Synonyms:	KCP2
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL4</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_173852, the custom clone sequence may differ by one or more nucleotides

```
ATGCGCATAGCTAACCGCACCCGGTTCAGCTCGCCTTTCTTGGCCAGAGGCGCCGGTTGGACTCACGGGC  
GGGGCATGATGGTGGTGGGTACGGGCACCTCGCTGGCGCTCTCCTCCCTCCTGTCCCTGCTCTTTGC  
TGGGATGCAGATGTACAGCCGTCAGCTGGCCTCCACCGAGTGGCTCACCATCCAGGGCGGCCTGCTTGGT  
TCGGGTCTCTTCGTGTTCTCGCTCACTGCCTTCAATAATCTGGAGAATCTGTCTTTGGCAAAGGATTCC  
AAGCAAAGATCTCCCTGAGATTCTCCTGTGCCTCCTGTTGGCTCTCTTGCATCTGGCCTCATCCACCG  
AGTCTGTGTACCACCTGCTTCATCTTCCATGGTGGTCTGTACTACATCAACAAGATCTCCTCCACC  
CTGTACCAGGCAGCAGCTCCAGTCTCACACCAGCCAAGGTCACAGGCAAGAGCAAGAAGAGAAACTGA
```



[View online »](#)

<b>5' Read Nucleotide Sequence:</b>	>OriGene 5' read for NM_173852 unedited NNNCCCGGGGANNNNCCCCCCCCCGNCTTTCAGAATTGTATACAATCATATAGGCG GCCGCGNAATTCGCACCAGCCGGTTCAGCTCGCCTTCTTGGCCAGAGGCGCCGGTTGG ACTCACGGGCGGGGCAATGATGGTGGTGGGTACGGGCACCTCGCTGGCGCTCTCCTCCCTC CTGTCCCTGCTGCTCTTGTGCGGATGCAGATGTACAGCCGTAGCTGGCCTCCACCGAG TGGCTCACCATCCAGGGCGGCCGCTTGGTTCGGGTCTCTTCGTGTTCTCGCTCACTGCC TTCAATAATCTGGAGAATCTTGTCTTTGGCAAAGGATTCCAAGCAAAGATCTTCCCTGAG ATTCTCCTGTGCCTCCTGTTGGCTCTCTTGCATCTGGCCTCATCCACCGAGTCTGTGTC ACCACCTGCTTCACTTCTCCATGGTTGGTCTGTACTACATCAACAAGATCTCCTCCACC CTGTACCAGGCAGCAGCTCCAGTCTCACACCAGCCAAGGTCACAGGCAAGAGCAAGAAG AGAAACTGACCTGAATGTTCAATAAAGTTGATTCTTTGTAAAAAAAAAAAAAAAAAACT CGACTCTAGATTGCGGCCGGTTCATAGCTGTTTCTGAACAGATCCCGGTGGCATCCC TGTGACCCCTCCCAGTGCCTCTCCTGGCCCTGGAAGTGGCACTCCAGTGCCACCAGC CTTGTCCTAATAAAATTAAGTTGCATCATTTTGTCTGACTAGGTGCTCTTATATATTA TGGGGTGAGGGGGTGGGTATGGAACCAGGGNGCAAGTTGGAAGACACCTGTAGGCTGC CGGGTCTATTGGGAACCGCTGNATGCAGTGCACAATCTGGCTCACTGAATCTCGCTCCT GGGTCAGCGATTCTCTGCTCN
<b>Restriction Sites:</b>	NotI-NotI
<b>ACCN:</b>	NM_173852
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>RefSeq:</b>	<a href="#">NM_173852.3</a> , <a href="#">NP_776251.1</a>
<b>RefSeq Size:</b>	577 bp
<b>RefSeq ORF:</b>	489 bp
<b>Locus ID:</b>	200185
<b>UniProt ID:</b>	<a href="#">Q8N6L1</a>
<b>Cytogenetics:</b>	1q22
<b>Protein Families:</b>	Transmembrane

**Gene Summary:**

Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity (PubMed:22467853). May be involved in N-glycosylation of APP (amyloid-beta precursor protein). Can modulate gamma-secretase cleavage of APP by enhancing endoproteolysis of PSEN1 (PubMed:21768116). [UniProtKB/Swiss-Prot Function]