

Product datasheet for **SC102000**

PSMD13 (AK092845) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PSMD13 (AK092845) Human Untagged Clone
Tag:	Tag Free
Symbol:	PSMD13
Synonyms:	HSPC027; p40.5; Rpn9; S11
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)



[View online »](#)

5' Read Nucleotide Sequence:	<p>>OriGene 5' read for AK092845 unedited</p> <pre>CCGAATTTGTAATACGACTCACTATAGGGNCGGCCGCGCAATTCGGCACGAGGCTCTTA GCTCAGACGACACCCTCCCACCCCACTGCCACCACCTGAGGCCTTCCCTGACCATCTGA ACTGAGATGAAGTAGCTGCCCTCTCCCGTGATGCAGTTGTTGCCTCATTGCATGTCGC TTCTTGTGTACACGCACATCTGTCTTCCCACTAGAACTTAAGCTTCGTCAGGAAGTGG CCCTTAGATTTCTGGTTTGCCGTGGTCACCCCTGGTCAACCCTGGCAAATTGTGGCCTCA AGTACTTGTTCGGGGATTGAACAATGGCTACTGTCAGAAACTCCTGTAAGGCATTTTGCT GTTATCCTTCTTATCTAAGCATTAAAAGCTTGTCTTAAACAAGATATATGTCTAATATTA GGAAACTTTTTAGTATGTGGGTAATAAAGATCTCTATTTTCAGAGCCAATATTGACA AAACATCCTTATCAGTTCTATTTTTATTTGGAAAAATAGTTTAAAAATAGTTAATAAAA TTTTTCCAGATGACTTTCACACGACCTGCCAATCACAGACAACTCACTTTGAAGAAAT TGCCAAAAGTGCTAAAATCACAGTGAATGAGGTACGGTCCCTAGGCTCAGGGTGTAGAG CAGCAAAGCTGCCACATGGAGGAGTCAAGGCTCTGTGTGAGCGTGTGCTCCCTAGACA GTAAAAATGACCGGAGGAGCNGCATCTGCTTCTACANAGCCATCTGGTCCCATGGGGG TGTGTCAGGCTATCTTTGTCTACACACGCTCCCTCTCTGCACAGGTGGAGCTTCTGGTG ATGAAAGCCCTTTTCGTGGNNGCTGTGAAAGGCAGTATANACGAAGTGGACAAACGAGTC ACATGACCCTGGTGCAGNCCCGAGTGTGNNNATTGCACAGTGATGTGTTGAAACCATN ATCCACTCTGTGATTTGAGGGGTGTGTCTATACGNCTAGTTCATTGGA</pre>
3' Read Nucleotide Sequence:	<p>>OriGene 3' read for AK092845 unedited</p> <pre>NGGTTGACTAAGNNANCGCGNCCCGCATTNANNGATCGATTTTTTTTTTTTTTTTTTTTT TTT GGGGTTTTTTTTAGGAACATTTCAAACCCCAACGGAAAAAATTTCCCTCCCGTGGAT GGGGATGGAAATCCCCCTCGGAGTACACAACTTAGGGGGCCCCCTGCCTGGCACGTTT TTTTTTCCCGTTTACCTAAAACCCCTGAAACCCCCCTTTGGGGGAAAAACCCCAA AAAAACCCCTCCCTGGGGCCTGGCTCCCATCAGGGACAGTCTGTTTTTAAAAACAAAA CAGTTTGTCTTTAAACAGGATCCCCACCCCCCAATTTAATGTCCACCGTTTGAAC ACCCCGTTTTTTGGCTGCAAACCCCTTTTTCAGGGGAGTCAAAGGAAACCCACGGGGA ATTCACATCCGGGCCCAAAACTCCAGGGGCTTTCATTCCCTTGATCTGAAATGGAAAT CCCGACCCAGGGACCTTAAAACCCCAAGGAAAAACAGGCCGAGCGGCCGGGACAACACC TGATGGGGGACTTTACATTTCCAGCTAACTTTTTTGGGAGCTTTTGTAAAAAGGACCT CTTAAGTGGGGAAATGGGACCGGACTTTCCCGTTAATTTTTTCCAGGAACCTTTAAAA AACTTGCTGAAAGAAAGCCGACAGGGCCGAGGAAATCCCCCGGATCGCCTTTCCCGC AATGTTAGAACACGTGGGCTTCAATCCAGAACCCGACGTTTCTGATCAAACCTCTATC TTAAGATTTAAAGGCCGTN</pre>
Restriction Sites:	NotI-NotI
ACCN:	AK092845
Insert Size:	2200 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [AK092845.1](#)

RefSeq Size: 2200 bp

RefSeq ORF: 2200 bp

Locus ID: 5719

Cytogenetics: 11p15.5

Protein Pathways: Proteasome

Gene Summary: The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a non-ATPase subunit of the 19S regulator. Two transcripts encoding different isoforms have been described. [provided by RefSeq, Jul 2008]