

Product datasheet for SC102102

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

POLR2J2 (NM_032958) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: POLR2|2 (NM 032958) Human Untagged Clone

Tag: Tag Free Symbol: POLR2J2

Synonyms: DNA directed RNA polymerase II polypeptide J-related; HRPB11B; MGC54043; MGC105050;

polymerase (RNA) II (DNA directed) polypeptide J2; RPB11, HRPB11B, MGC54043,

MGC105050; RPB11b1

Mammalian Cell

Selection:

None

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >NCBI ORF sequence for NM_032958, the custom clone sequence may differ by one or more

nucleotides

ATGAACGCCCCTCCAGCCTTCGAGTCGTTCTTGCTCTTCGAGGGCGAGAAGATCACCATTAACAAGGACA CCAAGGTACCCAAGGCCTGCTTATTCACCATCAACAAAGAAGACCACACACTGGGAAACATCATTAAATC ACGTGCCTGCTTCCCCTTCGCCTTCTGCCGTGATTGTCAGTTTCCTGAGGCCTCCCCAGCCACGCTTCCT GTACAGCCTGCAGAACTCTGCCCCAGAGCACATCAGCTATGTGCCCCAGCTCTCAAACGACACCTTGGCG GGGAGGCTCACCCTGTCCACCTTCACGCTGGAGCAGCCTCTAGGCCAGTTCAGCAGCCACAACATCTCTG

Α



5' Read Nucleotide Sequence: >OriGene 5' read for NM_032958 unedited

CGTCGCCATTTGTATACGACTCCTATAGGGCGGCCGCGAATTCGCACGAGGCTGGACGCA ACGGCGGCGGAGCATGAACGCCCCTCCAGCCTTCGAGTCGTTCTTGCTCTTCGAGGGCG AGAAGATCACCATTAACAAGGACACCAAGGTACCCAATGCCTGCTTATTCACCATCAACA AAGAAGACCACACACTGGGAAACATCATTAAATTACGTGCCTGCTTCCCCTTCGCCTTCT GCCGTGATTGTCAGTTTCCTGAGGCCTCCCCAGCCACGCTTCCTGTACAGCCTGCAGAAC TCTGCCCCAGAGCACATCAGCTATGTGCCCCAGCTCTCAAACGACACCTTGGCGGGGAGG CTCACCCTGTCCACCTTCACGCTGGAGCAGCCTCTAGGCCAGTTCAGCAGCCACAACATC TCTGACTTGGATACCATCTGGCTGGTGGTGGCCCTCAGCAACGCCACCCAGAGCTTCACG GCCCACGGACAAACCAGGACATCCCTGCTCCTGCCAACTTCTCCCAGAGGGGCTACTAT GTCGGCAATGATACCCACTGCCAACCAACAAAAATTGGCTGCAACCATCCCCTACCAGGA CCCGGCCCCTACAGGGTGAAGTTCCTGGTGATGAATGACGAAGGACCCGTGGCTGAAACA AATGGTCCAGCGACACTCGCCTGCAGCAAGCCCAGCACTTTGGGCTGTCCCCGGCCCCCA NAGCCCGGCACCGTGGTCATAATGGCCCTCCTGTTTATNCTCCTGGCCGGCCTCCTCACG GGACTCCTTGCCTGGGCTCATATACCCCTGCTTCAAAAGCTTGAGGAGCCCTTCCCTATC AGGCCCCCAGAAGG

Restriction Sites: Notl-Notl

ACCN: NM_032958

Insert Size: 3830 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 032958.3</u>, <u>NP 116580.2</u>

RefSeq Size: 1552 bp
RefSeq ORF: 351 bp
Locus ID: 246721
Cytogenetics: 7q22.1

Protein Families: Transcription Factors

Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase



Gene Summary:

This gene is a member of the RNA polymerase II subunit 11 gene family, which includes three genes in a cluster on chromosome 7q22.1 and a pseudogene on chromosome 7p13. The founding member of this family, DNA directed RNA polymerase II polypeptide J, has been shown to encode a subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. This locus produces multiple, alternatively spliced transcripts that potentially express isoforms with distinct C-termini compared to DNA directed RNA polymerase II polypeptide J. Most or all variants are spliced to include additional non-coding exons at the 3' end which makes them candidates for nonsense-mediated decay (NMD). Consequently, it is not known if this locus expresses a protein or proteins in vivo. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (2), also called beta, differs in the 3' coding region and UTR, compared to variant 1. Isoform 2 has a distinct C-terminus and is shorter than isoform 1.