OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for SC107411

LCLAT1 (NM_182551) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	LCLAT1 (NM_182551) Human Untagged Clone
Tag:	Tag Free
Symbol:	LCLAT1
Synonyms:	1AGPAT8; AGPAT8; ALCAT1; HSRG1849; LYCAT; UNQ1849
Mammalian Cell Selection:	None
Vector:	pCMV6-XL4
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_182551, the custom clone sequence may differ by one or more nucleotides

ATGCATTCCAGGGGAAGGGAAATTGTGGTGCTTCTGAACCCATGGTCAATTAACGAGGCAGTTTCTAGCT
ACTGCACGTACTTCATAAAGCAGGACTCTAAAAGCTTTGGAATCATGGTGTCATGGAAAGGGATTTACTT
TATACTGACTCTGTTTTGGGGAAGCTTTTTTGGAAGCATTTTCATGCTGAGTCCCTTTTTACCTTTGATG
TTTGTAAACCCATCTTGGTATCGCTGGATCAACAACCGCCTTGTGGCAACATGGCTCACCCTACCTGTGG
CATTATTGGAGACCATGTTTGGTGTAAAAGTGATTATAACTGGGGATGCATTTGTTCCTGGAGAAAGAAG
TGTCATTATCATGAACCATCGGACAAGAATGGACTGGATGTTCCTGTGGAATTGCCTGATGCGATATAGC
TACCTCAGATTGGAGAAAATTTGCCTCAAAGCGAGTCTCAAAGGTGTTCCTGGATTTGGTTGGGCCATGC
AGGCTGCTGCCTATATCTTCATTCATAGGAAATGGAAGGATGACAAGAGCCATTTCGAAGACATGATTGA
TTACTTTTGTGATATTCACGAACCACTTCAACTCCTCATATTCCCAGAAGGGACTGATCTCACAGAAAAC
AGCAAGTCTCGAAGTAATGCATTTGCTGAAAAAAATGGACTTCAGAAATATGAATATGTTTTACATCCAA
GAACTACAGGCTTTACTTTTGTGGTAGACCGTCTAAGAGAAGGTAAGAACCTTGATGCTGTCCATGATAT
CACTGTGGCGTATCCTCACAACATTCCTCAATCAGAGAAGCACCTCCTCCAAGGAGACTTTCCCAGGGAA
ATCCACTTTCACGTCCACCGGTATCCAATAGACACCCTCCCCACATCCAAGGAGGACCTTCAACTCTGGT
GCCACAAACGGTGGGAAGAGAAAGAAGAGAGGCTGCGTTCCTTCTATCAAGGGGAGAAGAATTTTTATTT
TACCGGACAGAGTGTCATTCCACCTTGCAAGTCTGAACTCAGGGTCCTTGTGGTCAAATTGCTCTCTATA
CTGTATTGGACCCTGTTCAGCCCTGCAATGTGCCTACTCATATATTTGTACAGTCTTGTTAAGTGGTATT
TTATAATCACCATTGTAATCTTTGTGCTGCAAGAGAGAATATTTGGTGGACTGGAGATCATAGAACTTGC
ATGTTACCGACTTTTACACAAACAGCCACATTTAAATTCAAAGAAAAATGAGTAA

View online »

5' Read Nucleotide Sequence:

>OriGene 5' read for NM_182551 unedited
AGGAAGGGGAGCTGCAGCCCGGGCTTTTCCTCCTGAGCCATGGACCTTCCCCGGGCCGCG
ACCTCGGGCTGAAAAACAGAGTGGGTACTCTCTTCTGGGAAGCTGGCAACAAATGGATGA
TGTGATATATGCATTCCAGGGGAAGGGAAATTGTGGTGCTTCTGAACCCATGGTCAATTA
ACGAGGCAGTTTCTAGCTACTGCACGTACTTCATAAAGCAGGACTCTAAAAGCTTTGGAA
TCATGGTGTCATGGAAAGGGATTTACTTTATACTGACTCTGTTTTGGGGAAGCTTTTTTG
GAAGCATTTTCATGCTGAGTCCCTTTTTACCTTTGATGTTTGTAAACCCATCTTGGTATC
GCTGGATCAACAACCGCCTTGTGGCAACATGGCTCACCCTACCTGTGGCATTATTGGAGA
CCATGTTTGGTGTAAAAGTGATTATAACTGGGGATGCATTTGTTCCTGGAGAAAGAAGTG
TCATTATCATGAACCATCGGACAAGAATGGCCTGGGATNGTTCCTTNGTGGAATTGCCTG
ATGCGATATAGCTACCTCAGATTGGAGAAAATTTGCCTCANAGCAGTTCAAAAAGGTGTT
CCTGGATTTTGGGTTGGGCCATGCAGGCTGCTGCCTATATCTTCATTCTANGAANAGGGN
AGGGATGACAGAGCCATTTCGAGACATGATGATAACTTTTTGTGATATTACGAACACTTC
AACTCTATATTCCAGAAGGGACTGATCTACAGAAACAGAGNNTCTCGAAGTATGCATTGC
TGAAAAAATGACTCGAATANTAA

3' Read Nucleotide Sequence:

>OriGene 3' read for NM_182551 unedited
NCCAAGACTTGNAACCGCGGCCGCATNCTANGATCGAGTTTTTTTTTTTTTTTTTTTTT
TTTTTTTTTTTAAAACTATTTAATTCACTCCTTTATTCTGGGATGTATATTACAGATAAC
ACAACTCACAAATATACCATCAGACATTGAAAACTAAGGCCATTCTGTGAGTTATTTTTA
AAACTTGGTGTTTTGCACATAATGATCTTAAAAAAAAATGAATTACCAAAACCAAGATTC
TCTTCTAAAATGAAAATTTAATGCAGGTACAGGATAACTTTAGGGCTATATCTAATCTGA
AGCTTATCAGGTAGCAAAACCATTTTCGTTTTCTACAGCATAAATAACAGCTCTAAGGCA
ACCACTACCTCAGCATGAAGCTCATTTCTCCACGTTAGAGTAGTGTTTACCTGCTACAGT
GACCAGTGTTTAAAGACCATTTCCCTTTCAGTAGCAAAAAGAGACTTTACCTAAGAAACA
CACTACATACTACAGAATCCTTGGAACAAGAAACAGAAAGGGAGCTGTAACTAAGGCACT
GAAAGCACATTATTTGTATAAAGAAATGTAAACAATTTAACACCAACAGGCTCCCTCCGT
TGGAGTCTTTATTGTAAATGCACACAGGGGCATCATAGTCAGGGCTAGAGTTTGAAGTCT
GGGTGAAACAAATGTTTCTGCAAAGTTCAAAGTCTTGCCTGCCAATATATTCTAACATGG
TGCTGTTGGAATAACTATTTCAAAGTGACCCAGTGGAAAGCTGGGATTCCCTTGAATCCG
AGTTTTAATAGCCTTACCAAACAGGAAGAAATTTTAGAGAAGTCCAAAATTTTGGGGCTC
TTTCAAACATTAATTTGATTCCTTTATAAGTTCCTTCAAGGGATTTTAGGATAATTTGGG
CTAATCCC

Restriction Sites: NotI-NotI

ACCN: NM_182551

Insert Size: 4500 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_182551.3, NP_872357.2
RefSeq Size:	5077 bp
RefSeq ORF:	1245 bp
Locus ID:	253558
UniProt ID:	Q6UWP7
Cytogenetics:	2p23.1
Protein Families:	Transmembrane
Protein Pathways:	Ether lipid metabolism, Glycerolipid metabolism, Glycerophospholipid metabolism, Limonene and pinene degradation, Metabolic pathways, Phenylalanine metabolism, Tyrosine metabolism
Gene Summary:	Exhibits acyl-CoA:lysocardiolipin acyltransferase (ALCAT) activity; catalyzes the reacylation of lyso-cardiolipin to cardiolipin (CL), a key step in CL remodeling (By similarity). Recognizes both monolysocardiolipin and dilysocardiolipin as substrates with a preference for linoleoyl-CoA and oleoyl-CoA as acyl donors (By similarity). Also exhibits 1-acyl-sn-glycerol-3-phosphate acyltransferase activity (AGPAT) activity; converts 1-acyl-sn-glycerol-3- phosphate (lysophosphatidic acid or LPA) into 1,2-diacyl-sn-glycerol-3- phosphate (phosphatidic acid or PA) by incorporating an acyl moiety at the sn-2 position of the glycerol backbone (PubMed:16620771). Possesses both lysophosphatidylinositol acyltransferase (LPIAT) and lysophosphatidylglycerol acyltransferase (LPGAT) activities (PubMed:19075029). Required for establishment of the hematopoietic and endothelial lineages (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (1) encodes the longest isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.