

Product datasheet for **SC108044**

CEACAM1 (NM_001712) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CEACAM1 (NM_001712) Human Untagged Clone
Tag:	Tag Free
Symbol:	CEACAM1
Synonyms:	BGP; BGP1; BGPI
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >SC108044 representing NM_001712.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGGAATTCGTGACTG
GATCCGGTACCGAGGAGATCTGCCGCCACGATCGCC
ATGGGGCACCTCTCAGCCCCACTTCACAGAGTGGTGTACCCTGGCAGGGGCTTCTGCTCACAGCCTCA
CTTCTAACCTTCTGGAACCCGCCACCCTGCCCAGCTCACTACTGAATCCATGCCATTCAATGTTGCA
GAGGGGAAGGAGGTTCTTCTCCTTGTCCACAATCTGCCCCAGCAACTTTTGGCTACAGCTGGTACAAA
GGGGAAGAGTGGATGGCAACCGTCAAATTGTAGGATATGCAATAGGAACTCAACAAGCTACCCAGGG
CCCGCAACAGCGGTGAGAGACAATATACCCCAATGCATCCCTGCTGATCCAGAACGTACCCAGAAT
GACACAGGATTCTACACCCTACAAGTCATAAAGTCAGATCTTGTGAATGAAGAAGCAACTGGACAGTTC
CATGTATACCCGGAGCTGCCAAGCCCTCCATCTCCAGCAACAACCTCAACCCCTGTGGAGGACAAGGAT
GCTGTGGCCTTCACTGTGAACCTGAGACTCAGGACACAACCTACCTGTGGTGGATAAACAATCAGAGC
CTCCCGGTGAGTCCAGGCTGCAGCTGTCCAATGGCAACAGGACCCTCACTCTACTCAGTGCACAAGG
AATGACACAGGACCCTATGAGTGTGAAATACAGAACCAGTGAGTGCGAACCGCAGTGACCCAGTCACC
TTGAATGTCACCTATGGCCCGGACACCCCAACATTTCCCTTCCAGACACCTATTACCGTCCAGGGGCA
AACCTCAGCCTCTCCTGCTATGCAGCCTTAACCCACCTGCACAGTACTCCTGGCTTATCAATGGAACA
TTCCAGCAAAGCACACAAGAGCTCTTTATCCCTAACATCACTGTGAATAATAGTGGATCCTATACCTGC
CACGCCAATAACTCAGTCACTGGCTGCAACAGGACCACAGTCAAGACGATCATAGTCACTGAGCTAAGT
CCAGTAGTAGCAAAGCCCCAAATCAAAGCCAGCAAGACCACAGTCAAGGAGATAAAGACTCTGTGAAC
CTGACCTGCTCCACAAATGACTGGAATCTCCATCCGTTGGTCTTCAAAAACAGAGTCTCCCGTCC
TCGGAGAGGATGAAGCTGTCCAGGGCAACACCACCTCAGCATAAACCCGTCAAGAGGGAGGATGCT
GGGAGTATTGGTGTGAGGTCTTCAACCCAATCAGTAAGAACCAAAGCGACCCCATCATGCTGAACGTA
AACTATAATGCTCTACCACAAGAAAATGGCCTCTCACCTGGGGCCATTGCTGGCATTGTGATTGGAGTA
GTGCCCTGTTGCTCTGATAGCAGTAGCCCTGGCATGTTTTCTGCATTTGCGGAAGACCGGCAGGGCA
AGCGACCAGCGTATCTCACAGAGCACAAACCCTCAGTCTCCAACCACACTCAGGACCACTCCAATGAC
CCACCTAACAGATGAATGAAGTTACTTATTCTACCCTGAACTTTGAAGCCAGCAACCCACACAACCA
ACTTCAGCCTCCCATCCCTAACAGCCACAGAAATAATTTATTTCAGAAGTAAAAAGCAGTAA
ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
  
```

Chromatograms: https://cdn.origene.com/chromatograms/ja3103_e10.zip

Restriction Sites: SgfI-MluI

ACCN: NM_001712

Insert Size: 1581 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001712.4
RefSeq Size:	3528 bp
RefSeq ORF:	1581 bp
Locus ID:	634
UniProt ID:	P13688
Cytogenetics:	19q13.2
Domains:	ig, IGc2, IG
Protein Families:	Druggable Genome, Transmembrane
MW:	57.6 kDa
Gene Summary:	<p>This gene encodes a member of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily. Two subgroups of the CEA family, the CEA cell adhesion molecules and the pregnancy-specific glycoproteins, are located within a 1.2 Mb cluster on the long arm of chromosome 19. Eleven pseudogenes of the CEA cell adhesion molecule subgroup are also found in the cluster. The encoded protein was originally described in bile ducts of liver as biliary glycoprotein. Subsequently, it was found to be a cell-cell adhesion molecule detected on leukocytes, epithelia, and endothelia. The encoded protein mediates cell adhesion via homophilic as well as heterophilic binding to other proteins of the subgroup. Multiple cellular activities have been attributed to the encoded protein, including roles in the differentiation and arrangement of tissue three-dimensional structure, angiogenesis, apoptosis, tumor suppression, metastasis, and the modulation of innate and adaptive immune responses. Multiple transcript variants encoding different isoforms have been reported, but the full-length nature of all variants has not been defined. [provided by RefSeq, May 2010]</p> <p>Transcript Variant: This variant (1) represents the longest transcript and encodes the longest protein (isoform 1). This variant has been referred to by multiple names, including transmembrane carcinoembryonic antigen BGP_a, TM1-CEA, and CEACAM1-4L. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>