

Product datasheet for **SC113612**

Transmembrane protein 30A (TMEM30A) (NM_018247) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Transmembrane protein 30A (TMEM30A) (NM_018247) Human Untagged Clone
Tag:	Tag Free
Symbol:	Transmembrane protein 30A
Synonyms:	C6orf67; CDC50A
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_018247, the custom clone sequence may differ by one or more nucleotides

```
ATGGCGATGAACTATAACGCGAAGGATGAAGTGGACGGTGGGCCCCCGTGTGCTCCGGGGGGCACCGCCA
AGACTCGGAGACCGGATAACACGGCCTTCAAACAGCAACGGCTGCCAGCTTGGCAGCCCATCCTTACGGC
TGGCACGGTGCTACCTATTTTCTTCATCATCGGTCTCATCTTCATTCCCATCGGCATTGGCATTTTTGTG
ACCTCCAACAACATCCGCGAGATCGAGATTGATTATACCGGAACAGAGCCTTCCAGTCCCTGTAATAAAT
GTTTATCTCCGGATGTGACACCTTGCTTTTGTACCATTAACCTCACACTGGAAAAGTCATTTGAGGGCAA
CGTGTATGTATTATGGACTGTCTAATTTCTATCAAAACCATCGTCGTTACGTGAAATCTCGAGATGAT
AGTCAACTAAATGGAGATTCTAGTGCTTTGCTTAATCCCAGTAAGGAATGTGAACCTTATCGAAGAAATG
AAGACAAACCAATTGCTCCTTGTGGAGCTATTGCCAACAGCATGTTTAATGATACATTAGAATTGTTTCT
CATTGGCAATGATTCTTATCCTATACCTATCGCTTTGAAAAAGAAAGGATTGCTTGGTGACAGATAAAA
AATGTGAAATTCAGAAATCCCCCTGGAGGAGACAACTGGAAGAACGATTAAAGGTACAACAAAGCCTG
TGAAGTGGCTTAAACCAGTTTACATGCTGGATTCTGACCCAGATAATAATGGATTCATAAATGAGGATTT
TATTGTTTGGATGCGTACTGCAGCATTACCTACTTTTCGCAAGTTGTATCGTCTTATAGAAAAGGAAAAGT
GATTTACATCCAACATTACCAGCTGGCCGATACTCTTTGAATGTCACATAACAATTACCCTGTACATTATT
TTGATGGACGAAAACGGATGATCTTGAGCACTATTTTATGGATGGGAGGAAAAAATCCATTTTGGGGAT
TGCTTACATCGCTGTTGGATCCATCTCCTTCTGGGAGTTGTACTGCTAGTAATTAATCATAAATAT
AGAAACAGTAGTAATACAGCTGACATTACCATTTAA
```



[View online »](#)

5' Read Nucleotide Sequence:

>OriGene 5' read for NM_018247 unedited
 GTTTANGATTTTGTAAATCCGACTTACTATAGGCGGCCGCGCAATTCGCACGAGGTGGCTG
 CGGCGCGCGCGCGCGCAGCGGCGCTCGAGCGGTTCTGTGAGGGTCAGCCGGCGGGCCCC
 CTGGGTGGTCCACCTGCAAAATCGCGGAGCGGCGCCCCAGGGATCGATGGCGATGAACTAT
 AACGCGAAGGATGAAGTGGACGGTGGGCCCCGTGTGCTCCGGGGGGCACCGCGAAGACT
 CGGAGACCGGATAACACGGCCTTCAAACAGCAACGGCTGCCAGCTTGGCAGCCCATCCTT
 ACGGCTGGCAGCGTGCTACCTATTTTCTCATCATCGGTCTCATCTTCATTCCCATCGGC
 ATTGGCATTTTTGTACCTCCAACAACATCCGCGAGATCGAGATTGATTATACCGGAACA
 GAGCCTTCCAGTCCCTGTAATAAATGTTTATCTCCGGATGTGACACCTTGCTTTTGTACC
 ATTAACCTCACACTGGAAAAGTCATTTGAGGGCAACGTGTTTATGTATTATGGACTGTCT
 AATTTCTATCAAACCATCGTCGTTACGTGAAATCTCGAGATGATAGTCAACTAAATGGA
 GATTCTAGTGCTTTGCTTAATCCCAGTAAGGAATGTGAACCTTATCGAAGAAATGAAGAC
 NAACCAATTGCTCCTTGTGGAGCTATTGCCAACAGCATGTTTAAATGATACATTAGAATTG
 TTTCTCATTGGCAATGATTCTTATCCTATACCTATCGCTTTTAAAAGAAAGGATTGCT
 TGGTGGACAGATAANAATGTNGAATTCCAGAAATCCNCTGGNNAGAGACCTGNNAGAA
 CGATTTANAGGGTACACANAGCCTGTGAACTGGNCTAACCAAGTTNACATGCNTGNATCTG
 ACCCCAGATATATGGATTATAAATGAG

3' Read Nucleotide Sequence:

>OriGene 3' read for NM_018247 unedited
 CGGCGCCCCGCCCCNNNNNNNNCCNNNNNGNNTGTCTTGGTGGCGTCCATTTTACC
 GTTAAATCTTATGAACACACTTCTTTAACTGATTTCTATGTTATATAAACTGAAAG
 TGTGGCACATTTCTACTATTAACCATTACATTGTTTTTGTTTTTAAAAAAGTATATA
 TTAATCAAAGTATACATTACCCAGACCTTTTGATTTAAATATAACTGACAGGAAAATA
 TAAACTCTGAGTAAACAGGCATTCTTCTACACAACATGAGTTATTTTCAGTAAAAAACA
 ACAACAAAAAGCAATTTCAATGTAATTAAGCCACCTCTGATTAATACATATTTTCATGC
 ATGTGGAAAAATTATTTTAACTATAACTTTAAGAATAAGTTCCATAGCAACAAGTTAC
 AGCATTATTTCAATAAGGAATATTAATAAATTCATTATTAATAAATAAATAAATTCAA
 CCATATACATATTATAATTTATCATGTACATACTCCAAAATCCCCATAATTTAATTAGGA
 AATTTATATAACATCTTTTTAAAGTATGTCTTAGGAAAACCAACCAAAACAATTGGCGCC
 TTAACTAATGAGAATGTGCATACACCATTGCTATTTGACCTATTATGCAAAAAGTGAATG
 AAATACATAGGAAATTAAGTTTCTGCTTTCAAACCTTTCTGGGAATATAGTAATTAATT
 AAAAACACTTACCTGCTGTAATAAAACAGCCTGAGCATTTCCAAAGTTGAAGTAATTTTC
 TTAATGTTAATCTCCCCATAATGAAAACCATTTCTTTTATTCCCTTTCTGTGGTTTACA
 CCTCCCGCAAAACACCACTGTTCCCATCAATTTCCCTAATACCCAACTCTTTCTTCC
 CCTA

Restriction Sites:

NotI-NotI

ACCN:

NM_018247

Insert Size:

2870 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_018247.2](#), [NP_060717.1](#)

RefSeq Size: 4410 bp

RefSeq ORF: 1086 bp

Locus ID: 55754

UniProt ID: [Q9NV96](#)

Cytogenetics: 6q14.1

Domains: CDC50

Protein Families: Druggable Genome, Transmembrane

Gene Summary:

Accessory component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids from the outer to the inner leaflet of various membranes and ensures the maintenance of asymmetric distribution of phospholipids. Phospholipid translocation seems also to be implicated in vesicle formation and in uptake of lipid signaling molecules. The beta subunit may assist in binding of the phospholipid substrate. Required for the proper folding, assembly and ER to Golgi exit of the ATP8A2:TMEM30A flippase complex. ATP8A2:TMEM30A may be involved in regulation of neurite outgrowth, and, reconstituted to liposomes, predominantly transports phosphatidylserine (PS) and to a lesser extent phosphatidylethanolamine (PE). The ATP8A1:TMEM30A flippase complex seems to play a role in regulation of cell migration probably involving flippase-mediated translocation of phosphatidylethanolamine (PE) at the plasma membrane. Required for the formation of the ATP8A2, ATP8B1 and ATP8B2 P-type ATPase intermediate phosphoenzymes. Involved in uptake of platelet-activating factor (PAF), synthetic drug alkylphospholipid edelfosine, and, probably in association with ATP8B1, of perfosine. Also mediate the export of alpha subunits ATP8A1, ATP8B1, ATP8B2, ATP8B4, ATP10A, ATP10B, ATP10D, ATP11A, ATP11B and ATP11C from the ER to other membrane localizations.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data because no single transcript was available for the full length of the gene. The extent of this transcript is supported by transcript alignments.