

Product datasheet for **SC116100**

TIP30 (HTATIP2) (NM_006410) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	TIP30 (HTATIP2) (NM_006410) Human Untagged Clone
Tag:	Tag Free
Symbol:	TIP30
Synonyms:	CC3; SDR44U1; TIP30
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL4</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>OriGene ORF within SC116100 sequence for NM_006410 edited (data generated by NextGen Sequencing) <pre> ATGGCCGAAACAGAACCCCTGTGCAAGCTTCGGGAAGACTTCAGGATGCAGAATAAATCC GTCTTTATTTTGGCGCCAGCGGAGAAACCGGCAGAGTGCTCTTAAAGGAAATCCTGGAG CAGGGCCTGTTTTCCAAAGTCACGCTCATTGGCCGGAGGAAGCTCACCTTCGACGAGGAA GCTTATAAAATGTGAATCAAGAAGTGGTGGACTTTGAAAAGTTGGATGACTACGCCTCT GCCTTTCAAGGTCATGATGTTGGATTCTGTTGCCTGGGTACCACCAGAGGAAAGCTGGG GCGGAGGGATTGTTCTGTGTTGACCGAGATTATGTGCTGAAGTCTGCAGAGCTGGCAAAA GCTGGAGGGTGCAAACATTTCAACTTGCTATCCTCTAAAGGAGCTGATAAATCAAGCAAT TTTTTATATCTACAAGTTAAGGGAGAAGTAGAAGCCAAGGTTGAAGAATTAATTTGAT CGTTACTCTGTATTTAGGCCTGGAGTTCTGTTATGTGATAGGCAAGAATCTCGCCAGGT GAATGGCTGGTTAGAAAGTTCTTTGGCTCCTTACCAGACTCTTGGGCCAGTGGGCATTCT GTGCCTGTGGTGACCGTGGTTAGAGCAATGCTGAACAATGTGGTGAGACCAAGAGACAAG CAGATGGAAGTCTGGAGAAACAAGGCCATCCATGACCTGGGGAAGCGCATGGCTCTCTC AAGCCATGA </pre>

Clone variation with respect to NM_006410.4



[View online »](#)

5' Read Nucleotide Sequence:	>OriGene 5' read for NM_006410 unedited <pre> TTAGATTTTGTAAACAGCTCACTATAGGGCGGCCGCTCTGATGGCCGGGCTGCGGC GCTGAGCGCGGGCGGGCGGCTGCTCTGGCGGCCGCCCTGCTCCTGCTGCGTCTGAGGA CCCGGGGCGGGGGCTGGCCCCAGCATGGCCGAAACAGAAGCCCTGTGGAAGCTTCGGGA AGACTTCAGGATGCAGATAAATCCGTCTTTATTTGGGCGCCAGCGGAGAAACCGGCAG AGTGCTCTTAAAGGAAATCCTGGAGCAGGGCCTGTTTTCAAAGTCACGCTCATTGGCCG GAGGAAGCTCACCTTCGACGAGGAAGCTTATAAAATGTGAATCAAGAAGTGGTGGACTT TGAAAAGTTGGATGACTACGCCCTCTGCCTTTCAAGGTCATGATGTTGGATTCTGTTGCCT GGGTACCACCAGAGGAAAGCTGGGGCGGAGGGATTTGTTCTGTGTTGACCGAGATTATGT GCTGAAGTCTGCAGAGCTGGCAAAGCTGGAGGGTGCAAACATTTCAACTTGCTATCCTC TAAAGGAGCTGATAAATCAAGCAATTTTTATATCTACAAGTTAAGGGAGAAGTAGAAGC CAAGGTTGAAGAATTAATTTGATCGTTACTCTGTATTTAGGCCTGGAGTTCTGTTATG TGATAGGCAAGAATCTCGCCAGGTGAATGGCTGGTTAGAAAGTTCTTTGGCTCCTTACC AGACTCTTGGGCCAGTGGGCATTCTGTGCCTGTGGTGACCGTGGTTAGAGCAATGCTGAA CAATGTGGTGAGACCAAGAGACAGCAGATGGAAGTCTGGAGAACCAAGCCATCCTGACCT GGGAAGCGCATGGCTCTCTCAGCCATGACCACATGGAGAATGTTTTTTATTGTCACTTA CACCATCACAATTCGTATTCAGGTCTAAAAAGTACATGTTTACTTTTGTTACATC </pre>
Restriction Sites:	NotI-NotI
ACCN:	NM_006410
Insert Size:	1650 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_006410.3 , NP_006401.2
RefSeq Size:	1438 bp
RefSeq ORF:	729 bp
Locus ID:	10553
UniProt ID:	Q9BUP3
Cytogenetics:	11p15.1
Protein Families:	Druggable Genome

Gene Summary:

Oxidoreductase required for tumor suppression. NADPH-bound form inhibits nuclear import by competing with nuclear import substrates for binding to a subset of nuclear transport receptors. May act as a redox sensor linked to transcription through regulation of nuclear import. Isoform 1 is a metastasis suppressor with proapoptotic as well as antiangiogenic properties. Isoform 2 has an antiapoptotic effect.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) uses an alternate 5' exon and differs in the 5' UTR, compared to variant 1. These differences cause translation initiation at a downstream AUG and an isoform (b, also known as CC3) with a shorter N-terminus compared to isoform a. Variants 2, 3 and 4 encode the same isoform.