

Product datasheet for SC116227

PI 3 Kinase catalytic subunit alpha (PIK3CA) (NM_006218) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	PI 3 Kinase catalytic subunit alpha (PIK3CA) (NM_006218) Human Untagged Clone
Tag:	Tag Free
Symbol:	PI 3 Kinase catalytic subunit alpha
Synonyms:	CLAPO; CLOVE; CWS5; MCAP; MCM; MCMTC; p110-alpha; PI3K; PI3K-alpha
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>OriGene ORF within SC116227 sequence for NM_006218 edited (data generated by NextGen Sequencing)

```

ATGCCTCCACGACCATCATCAGGTGAAGTGTGGGCATCCACTTGATGCCCCCAAGAATCCTAGTAGAAT
GTTTACTACCAAATGGAATGATAGTGACTTTAGAATGCCTCCGTGAGGCTACATTAATAACCATAAAGCA
TGAAGTATTTAAAGAAGCAAGAAAATACCCCTCCATCAACTTCTCAAGATGAATCTTCTTACATTTTC
GTAAGTGTACTCAAGAAGCAGAAAGGGAAGAATTTTTGATGAAACAAGACGACTTTGTGACCTTCGGC
TTTTTCAACCCTTTTTAAAAGTAATTGAACAGTAGGCAACCGTGAAGAAAAGATCCTCAATCGAGAAAT
TGGTTTTGCTATCGGCATGCCAGTGTGTAATTTGATATGGTTAAAGATCCAGAAGTACAGGACTCCGA
AGAAATATTCTGAACGTTTGTAAAGAAGCTGTGGATCTTAGGGACCTCAATTCACCTCATAGTAGAGCAA
TGTATGTCTATCCTCAAATGTAGAATCTCACCAGAATTGCCAAAGCACATATAATAAATTAGATAA
AGGGCAAATAATAGTGGTATCTGGGTAATAGTTTCTCAAATAATGACAAGCAGAAGTATACTCTGAAA
ATCAACCATGACTGTGTACCAGAACAAGTAATTGCTGAAGCAATCAGGAAAAAACTCGAAGTATGTTGC
TATCCTCTGAACAATAAACTCTGTGTTTTAGAATATCAGGGCAAGTATATTTAAAAGTGTGTGGATG
TGATGAATACTTCTAGAAAAATATCCTCTGAGTCAGTATAAGTATAAAGAAGCTGATAATGCTTGGG
AGGATGCCCAATTTGATGTTGATGGCTAAAGAAAGCCTTATTCTCAACTGCCAATGGACTGTTTTACAA
TGCCATCTTATCCAGACGCATTTCCACAGCTACACCATATATGAATGGAGAAACATCTACAAAAATCCCT
TTGGGTTATAAATAGTGCACTCAGAATAAAAATCTTTGTGCAACCTACGTGAATGTAATATTCGAGAC
ATTGATAAGATCTATGTTGGAACAGGTATCTACCATGGAGGAGAACCCTTATGTGACAATGTGAACACTC
AAAGAGTACCTTGTTCATCCAGGTGGAATGAATGGCTGAATTATGATATATACATTCCTGATCTTCC
TCGTGCTGCTCGACTTTGCCCTTCCATTTGCTCTGTTAAAGGCCGAAAGGGTGCTAAAGAGGAACACTGT
CCATTGGCATGGGAAATATAAACTTGTGTTGATTACACAGACTCTAGTATCTGGAAAAATGGCTTTGA
ATCTTTGGCCAGTACCTCATGGATTAGAAGATTGCTGAACCCTATTGGTGTACTGGATCAAATCCAAA
TAAAGAACTCCATGCTTAGAGTTGGAGTTTACTGGTTCAGCAGTGTGGTAAAGTTCCAGATATGTCA
GTGATTGAAGAGCATGCCAATTTGGTCTGTATCCCGAGAAGCAGGATTTAGCTATCCACGCAGGACTGA
GTAACAGACTAGTAGAGACAATGAATTAAGGGAAAATGACAAAGAAGCAGCTCAAAGCAATTTCTACAGC

```



[View online »](#)

```

AGATCCTCTCTCTGAAATCACTGAGCAGGAGAAAGATTTTCTATGGAGTCACAGACACTATTGTGTAAC
ATCCCCGAAATCTACCCAAATTGCTTCTGTCTGTTAAATGGAATTCTAGAGATGAAGTAGCCAGATGT
ATTGCTTGGTAAAAGATTGGCCTCCAATCAAACCTGAACAGGCTATGGAACCTTCTGGACTGTAATTACCC
AGATCCTATGGTTCGAGGTTTTGCTGTTTCGGTCTTGGAAAAATATTTAACAGATGACAACTTTCTCAG
TATTTAATTCAGCTAGTACAGGTCCTAAAATATGAACAATATTTGGATAACTTGTGTGAGATTTTAC
TGAAGAAAAGCATTGACTAATCAAAGGATTGGGCACCTTTTCTTTTGGCATTAAAAATCTGAGATGCACAA
TAAAACGATTAGCCAGAGGTTTTGGCCTGCTTTTGGAGTCTATTGTCTGTCATGTGGGATGATTTGAAG
CACCTGAATAGGCAAGTCGAGGCAATGGAAAAGCTCATTAACTTAACTGACATTCTCAAACAGGAGAAGA
AGGATGAAAACACAAAAGGTACAGATGAAGTTTTAGTTGAGCAAATGAGGCGACCAGATTTTCATGGATGC
TCTACAGGGCTTTCTGTCTCCTCTAAACCTGCTCATCAACTAGGAAACCTCAGGCTTGAAGAGTGTGCA
ATTATGTCCTCTGCAAAAAGGCCACTGTGGTTGAATTGGGAGAACCAGACATCATGTCAGAGTTACTGT
TTCAGAACAATGAGATCATCTTTAAAAATGGGGATGATTTACGGCAAGATATGCTAACACTTCAAATAT
TCGTATTATGAAAATATCTGGCAAAATCAAGGCTTGATCTTCAATGTTACCTTATGGTTGTCTGTCA
ATCGGTGACTGTGTGGGACTTATTGAGGTGGTGCGAAATTTCTCACACTATTATGCAAATTCAGTGCAAAG
GCGGCTTGAAGGTGCACTGCAGTTCAACAGCCACACACTACATCAGTGGCTCAAAGACAAGAACAAGG
AGAAATATATGATGCAGCCATTGACCTGTTTACAGGTTTATGTGCTGGATCTGTGTAGCTACCTTCATT
TTGGGAATTGGAGATCGTCACAATAGTAACATCATGGTGAAGACGATGGACAACGTGTTTCATATAGATT
TTGGACACTTTTTGGATCACAAGAAGAAAAATTTGGTTATAAACGAGAACGTGTGCCATTTGTTTTGAC
ACAGGATTTCTTAATAGTGATTAGTAAAGGAGCCCAAGAATGCACAAAGACAAGAGAATTTGAGAGGTTT
CAGGAGATGTGTTACAAGGCTTATCTAGCTATTCGACAGCATGCCAATCTTTCATAAATCTTTTCTCAA
TGATGCTTGGCTCTGGAATGCCAGAACTACAATCTTTGATGACATTGCATACATTGAAAGACCCTAGC
CTTAGATAAAACTGAGCAAGAGGCTTTGGAGTATTTTATGAAACAATGAATGATGCACATCATGGTGGC
TGGACAACAAAAATGGATTGGATCTTCCACACAATTAACAGCATGCATTGAATGA
    
```

Clone variation with respect to NM_006218.3

5' Read Nucleotide Sequence:

```

>OriGene 5' read for NM_006218 unedited
NCCTGTTCAAATTTGTATCCGACTCACTATAGGCGGCCGCGCAATTCGCACGAGGAGGGA
GCGCACAAGAGGCTTTCTAGAAACAGTCTGCATATTCTTTAATGAATTAGAGTTGTGAT
TTCATCTTGGTGAAGCTTCTCATTCCAGCACTGCGCCTTTGTGCACATGCAGGACCTGG
GTTTTCGGAACGATGTTTTGAATCATAGTTTGTGTGGCCGGGAGAACCCTTAGAATAC
TTGGGTTTCTGCTTTGGGACAACCATACATCTAATTCCTTAAAGTAGTTTTATATGTAAA
ACTTGCAAAGAATCAGAACAATGCCTCCACGACCATCATCAGGTGAACGTGTTGGGCATCC
ACTTGATGCCCCAAGAATCCTAGTAGAATGTTTACTACCAAATGGAATGATAGTGACTT
TAGAATGCCTCCGTGAGGCTACATTAATAACCATAAAGCATGAACATTTAAAGAAGCAA
GAAAATACCCCTCCATCAACTTCTTCAAGATGAATCTTCTTACATTTTCGTAAGTGTTA
CTCAAGAAGCAGAAAGGGAAGAATTTTTGATGAAACAAGACGACTTTGTGACCTTCGGC
TTTTTCAACCCTTTTTAAAGTAATTGAACCAGTAGGCAACCGTGAAGAAAAGATCCTCA
ATCGAGAAATTTGTTTTGCTATCGGCATGCCAGTGTGTAATTTGATATGGTTAAAGATC
CAGAAGTACAGGACTTTGAAAGAATATTCTGAACGTTTGTAAAGAAGCTGTGGATCTTTT
AGGACCTCATTACCTCATAGTAGAGCAATGTATGTCTATCCTCCAATGTAGAATCTTCA
CCAGAATGCCAAAGCCATATTTTATAATAGATAAAGGCAATN
    
```

3' Read Nucleotide Sequence:	<p>>OriGene 3' read for NM_006218 unedited</p> <pre> ACCGCGGGCCGCAATCTANAGTCGAGTTTTTTTTTTTTTTTTTTTACTTTGTGGGAGCCCA GAATTTCCCTGGGAAATAAGAATAAGATCCTTTAAATACTGTCCAGAAGAAAAAAAAAAAA TACCACCTTATAACAGGTCTCATTGTGACTGCTTCCAAAAGTGCAGATAAAACCTCAA TCATCAAAGGTTTGATTTATTAAGAAAGAAAGTCTCTGGTGAATCTCAAGAGTCCCT TCCACCCCAAGCATTTCCTTAAACAAGCAGTCAAAAACCCACTTCAAACATAAAA TTGAACAGCAATGGAAAAATAAGAAATTTTCATTCTTTTCCAATCAATGTAATTTAACAT AAACTCCAGTTTACTTACACCTAATGCAATAACAGCCTTTGTTGTGCCACATTTCAAAA CATAGAACATTTGCAAGCAAAAAAACCATCACTTTTCTTCTCCATCATTTCTATATA TTTTGGGATTTTGTGTTTGTGTTTTTAAAAAAGATACAAACAATCTTCAAAGTT TACCTTTTGGACTTAAGGCATAACATGAAATTGCGCATTATTCTAAAGCTTAATTTTGA AATGAAGTAAAGTCTATCAAACCCTGTTTGCCTTACATTATTTAAATTATATAG NATTTTATTTCTGTTCTGCTGNAATCTAATGCTGTTTCAATGTTGCAATTCCTATG CNATCGGTCTTGCCTGCTGAGAGTTATTAACAGNTGCAGTGTGGAATCCAGAGTGAGCT TTCATTTTCTCAGNTATCTTTTTCAGTCATGCATGCTGTTAATGTGTGGAAGATCCATC CATTTTGTGTCAGCACATGATGTGCTCATTCAATNGNTATGAATACTNCAAAGCCTCT GCTCAGTTTACTAGGCTAAGGTCTTCAATGATGCATGCATAA </pre>
Restriction Sites:	NotI-NotI
ACCN:	NM_006218
Insert Size:	4530 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_006218.2 , NP_006209.2
RefSeq Size:	9104 bp

RefSeq ORF:	3207 bp
Locus ID:	5290
UniProt ID:	P42336
Cytogenetics:	3q26.32
Domains:	PI3K_rbd, PI3_PI4_kinase, PI3Ka, PI3K_C2, PI3K_p85B
Protein Families:	Druggable Genome
Protein Pathways:	Acute myeloid leukemia, Apoptosis, B cell receptor signaling pathway, Chemokine signaling pathway, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, ErbB signaling pathway, Fc epsilon RI signaling pathway, Fc gamma R-mediated phagocytosis, Focal adhesion, Glioma, Inositol phosphate metabolism, Insulin signaling pathway, Jak-STAT signaling pathway, Leukocyte transendothelial migration, Melanoma, mTOR signaling pathway, Natural killer cell mediated cytotoxicity, Neurotrophin signaling pathway, Non-small cell lung cancer, Pancreatic cancer, Pathways in cancer, Phosphatidylinositol signaling system, Progesterone-mediated oocyte maturation, Prostate cancer, Regulation of actin cytoskeleton, Renal cell carcinoma, Small cell lung cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway, Type II diabetes mellitus, VEGF signaling pathway
Gene Summary:	Phosphatidylinositol 3-kinase is composed of an 85 kDa regulatory subunit and a 110 kDa catalytic subunit. The protein encoded by this gene represents the catalytic subunit, which uses ATP to phosphorylate PtdIns, PtdIns4P and PtdIns(4,5)P2. This gene has been found to be oncogenic and has been implicated in cervical cancers. A pseudogene of this gene has been defined on chromosome 22. [provided by RefSeq, Apr 2016]