

Product datasheet for **SC118929**

CD89 (FCAR) (NM_002000) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	CD89 (FCAR) (NM_002000) Human Untagged Clone
Tag:	Tag Free
Symbol:	CD89
Synonyms:	CD89; CTB-61M7.2; FcalphaRI
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL4</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>OriGene ORF within SC118929 sequence for NM_002000 edited (data generated by NextGen Sequencing)

```

ATGGACCCCAAACAGACCACCCTCTGTGTCTTGTGCTCTGTCTGGGCCAGAGGATTCAG
GCACAGGAAGGGGACTTTCCCATGCCTTTCATATCTGCCAAATCGAGTCCTGTGATTCCC
TTGGATGGATCTGTGAAAATCCAGTGCCAGGCCATTTCGTGAAGCTTACCTGACCCAGCTG
ATGATCATAAAAACTCCACGTACCGAGAGATAGGCAGAAGACTGAAGTTTTGGAATGAG
ACTGATCCTGAGTTCGTTCATTGACCACATGGACGCAAACAAGGCAGGGCGCTATCAGTGC
CAATATAGGATAGGGCACTACAGATTCCGGTACAGTGACACCCTGGAGCTGGTAGTGACA
GGCTTGTATGGCAAACCTTCTCTCTGCAGATCGGGGTCTGGTGTGATGCCAGGAGAG
AATATTTCCCTCACGTGCAGCTCAGCACACATCCATTTGATAGATTTTCACTGGCCAAG
GAGGGAGAAGTTTCTCTGCCACAGCACCAAAGTGGGGAACACCCGGCCAAGTTCTCTTTG
GGTCCTGTGGACCTCAATGTCTCAGGGATCTACAGGTGCTACGGTTGGTACAACAGGAGC
CCCTACCTGTGGTCCCTTCCCCAGTAATGCCTTGGAGCTTGTGGTACAGACTCCATCCAC
CAAGATTACACGACGCAGAACTTGATCCGCATGGCCGTGGCAGGACTGGTCCCTCGTGGCT
CTCTTGGCCATACTGGTTGAAAATTGGCACAGCCATACGGCACTGAACAAGGAAGCCTCG
GCAGATGTGGCTGAACCGAGCTGGAGCCAACAGATGTGTCAGCCAGGATTGACCTTTGCA
CGAACACCAAGTGTCTGCAAGTAA

```

Clone variation with respect to NM_002000.2



[View online »](#)

5' Read Nucleotide Sequence:	>OriGene 5' read for NM_002000 unedited CAGAAATTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGCACGACACGGGGCTGA GGCCGTGTGACGACGATGGACCCCAAACAGACCACCTCTGTGTCTTGTGCTCTGTCTG GGCCAGAGGATTACGGACAGGAAGGGGACTTTCCCATGCCTTTTCATATCTGCCAAATCG AGTCTGTGATTCCTTGGATGGATCTGTGAAAATCCAGTGCCAGGCCATTCGTGAAGCT TACCTGACCCAGCTGATGATCATAAAAACTCCACGTACCGAGAGATAGGCAGAAGACTG AAGTTTTGGAATGAGACTGATCCTGAGTTCGTCATTGACCACATGGACGCAAACAAGGCA GGGCGCTATCAGTGCCAATATAGGATAGGGCACTACAGGTTCCGGTACAGTGACACCCTG GAGCTGGTAGTGACAGGCTTGTATGGCAAACCCTTCTCTCTGCAGATCGGGGTCTGGTG TTGATGCCAGGAGAGAATATTTCCCTCACGTGCAGCTCAGCACACATCCCATTTGATAGA TTTTCACTGGCCAAGGAGGGAGAACTTTCTCTGCCACAGCACAAAGTGGGGAACACCCG GCCAATTCTCTTTGGTCTGTGGACCTCAATGTCTCAGGGATCTACAGGTGCTACGGT TGGTACAACAGGAGCCCTACCTGTGGTCTTCCCAAGTAAATGCCTTGGAGCTTGTGGTC ACAGACTCCATCCACCAAGATTACACGACGAGAATTGATCCGCATGGCCGTGGCANGA CTGGTCTCGTGGCTCTTGGCATACTGGNTGAAAATGGCACAGNCATACGCACTGAAC AAGAAGCCTCGNCAGATGTGGCTGAACCGAGCTGGAGCACCAGATGTGTACGCCAGAATT GACCTTNGCACGACCACNNAGTGTCTGCAGTAACACCTGGGNAGTGAANGCAAANNAGA CCCAGACTGTTGNAGTCCGACAAGCTACTTGAAGACA
Restriction Sites:	ECoRI-NOT
ACCN:	NM_002000
Insert Size:	2620 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_002000.2</u> , <u>NP_001991.1</u>
RefSeq Size:	1671 bp
RefSeq ORF:	864 bp
Locus ID:	2204
UniProt ID:	<u>P24071</u>
Cytogenetics:	19q13.42
Domains:	ig, IG

Protein Families: Transmembrane

Gene Summary: This gene is a member of the immunoglobulin gene superfamily and encodes a receptor for the Fc region of IgA. The receptor is a transmembrane glycoprotein present on the surface of myeloid lineage cells such as neutrophils, monocytes, macrophages, and eosinophils, where it mediates immunologic responses to pathogens. It interacts with IgA-opsionized targets and triggers several immunologic defense processes, including phagocytosis, antibody-dependent cell-mediated cytotoxicity, and stimulation of the release of inflammatory mediators. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (1, also known as a.1) represents the longest transcript and encodes the longest isoform (a). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.