

Product datasheet for SC124566

HMGA1 (NM 145901) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: HMGA1 (NM_145901) Human Untagged Clone

Tag: Tag Free
Symbol: HMGA1

Synonyms: HMG-R; HMGA1A; HMGIY

Vector: pCMV6-XL6

E. coli Selection: Ampicillin (100 ug/mL)

Cell Selection: None

Fully Sequenced ORF: >NCBI ORF sequence for NM_145901, the custom clone sequence may differ by one or more

nucleotides

AAGAGGAGGCATCTCGCAGGAGTCCTCGGAGGAGGAGCAGTGA

5' Read Nucleotide Sequence: >OriGene 5' read for NM_145901 unedited

NNGGGCCATATCCCCCGCCCGTTGCCGCTTTGGGCGGTAGGCGTGTACGGTGGGAGGTCT

CTACTCCACTGCCTGGCACAGCAGGTGGGCCATGGAAGGGG

Restriction Sites: Notl-Notl



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORÏGENE

ACCN: NM_145901

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 145901.1, NP 665908.1</u>

 RefSeq Size:
 1876 bp

 RefSeq ORF:
 324 bp

 Locus ID:
 3159

 UniProt ID:
 P17096

 Cytogenetics:
 6p21.31

Protein Families: Druggable Genome, Stem cell - Pluripotency, Stem cell relevant signaling - JAK/STAT signaling

pathway, Transcription Factors

Gene Summary: This gene encodes a chromatin-associated protein involved in the regulation of gene

transcription, integration of retroviruses into chromosomes, and the metastatic progression of cancer cells. The encoded protein preferentially binds to the minor groove of AT-rich regions in double-stranded DNA. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been identified on multiple

chromosomes. [provided by RefSeq, Jan 2016]

Transcript Variant: This variant (3) differs in the 5' UTR compared to variant 1. Variants 1, 3, 9, 10, 12, and 13 encode isoform (a, also called HMG-I). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record

were based on transcript alignments.