

Product datasheet for **SC126413**

HIST2H2AC (BC060324) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	HIST2H2AC (BC060324) Human Untagged Clone
Tag:	Tag Free
Symbol:	HIST2H2AC
Synonyms:	H2A; H2A-GL101; H2A/q; H2AFQ; H2A histone family, member Q; histone 2, H2ac; histone cluster 2, H2ac; histone IIa; MGC74460
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Cell Selection:	None
Fully Sequenced ORF:	>OriGene sequence for BC060324 edited CTGAGCGTTGTCTGTGTTTAACTTGATTTTCAGTCATGTCTGGTCTGGCAAACAAGGAG GCAAGGCCCGCGCCAAGGCCAAGTCGCGCTCGTCCCGCGCTGGCCTCCAGTTCCCGGTAG GGCGAGTGCACCGCTTGTCTGCGCAAAGGCAACTACGCGGAGCGGGTGGGGCCGGCGCGC CCGTCTACATGGCGGCGGTCTCGAGTACCTGACCGCGGAGATCCTGGAGCTGGCGGGCA ACGCGGCTCGGGACAACAAGAAGACGCGCATCATCCCTCGTCACCTCCAGCTGGCCATCC GCAACGACGAGGAACTGAACAAGCTGCTGGCAAAGTCACCATCGCCCAGGGCGGCGTTT TGCCTAACATCCAGGCCGTTCTGTTACCAAAGAAAACCGAAAGCCACAAAGCCAAAAGCA AATAAATGCAGAAAAAAAAAAAAAAAAA
Restriction Sites:	NotI-NotI
ACCN:	BC060324
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC060324.1](#), [AAH60324.1](#)

RefSeq Size: 446 bp

Locus ID: 8338

Cytogenetics: 1q21.2

Protein Pathways: Systemic lupus erythematosus

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. [provided by RefSeq, Aug 2015]