

Product datasheet for SC126913

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

IL32 (NM_004221) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: IL32 (NM_004221) Human Untagged Clone

Tag: Tag Free Symbol: IL32

Synonyms: IL-32alpha; IL-32beta; IL-32delta; IL-32gamma; NK4; TAIF; TAIFa; TAIFb; TAIFc

Mammalian Cell None

Selection:

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene ORF within SC126913 sequence for NM_004221 edited (data generated by NextGen

Sequencing)

TGCTCTGAACCCCAATCCTCAAAATGA

Clone variation with respect to NM_004221.4

Restriction Sites: Notl-Notl
ACCN: NM 004221

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 004221.4</u>, <u>NP 004212.4</u>

 RefSeq Size:
 1090 bp

 RefSeq ORF:
 567 bp

 Locus ID:
 9235

 UniProt ID:
 P24001

 Cytogenetics:
 16p13.3

Protein Families: Secreted Protein

Gene Summary: This gene encodes a member of the cytokine family. The protein contains a tyrosine sulfation

site, 3 potential N-myristoylation sites, multiple putative phosphorylation sites, and an RGD cell-attachment sequence. Expression of this protein is increased after the activation of T-cells by mitogens or the activation of NK cells by IL-2. This protein induces the production of TNFalpha from macrophage cells. Alternate transcriptional splice variants, encoding different

isoforms, have been characterized. [provided by RefSeq, Jul 2008]

Transcript Variant: This variant (2) lacks an alternate exon in the 5' UTR compared to variant 1. Variants 1 and 2 encode the same protein, isoform B, also referred to as IL-32beta. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for

the transcript record were based on alignments.