

Product datasheet for **SC127163**

G protein beta subunit like (MLST8) (NM_022372) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	G protein beta subunit like (MLST8) (NM_022372) Human Untagged Clone
Tag:	Tag Free
Symbol:	G protein beta subunit like
Synonyms:	GbetaL; GBL; LST8; POP3; WAT1
Mammalian Cell Selection:	None
Vector:	<u>pCMV6-XL5</u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_022372, the custom clone sequence may differ by one or more nucleotides

```
ATGAACACCTCCCAGGCACGGTGGGCAGTGACCCGGTCATCCTGGCCACTGCAGGCTACGACCACCCG
TGCGCTTCTGGCAGGCCACAGCGGCATCTGCACCCGGACGGTGCAGCACCAGGACTCCCAGGTGAATGC
CTTGGAGGTCACACCGGACCGCAGCATGATTGCTGCTGCAGGTTACCAGCACATCCGCATGTATGATCTC
AACTCCAATAACCCCTAACCCCATCATCAGCTACGACGGCGTCAACAAGAACATCGCGTCTGTGGGCTTCC
ACGAAGACGGCCGCTGGATGTACACGGGCGGCGAGGACTGCACAGCCAGGATCTGGGACCTCAGTCCC
GAACCTGCAGTGCCAGCGGATCTCCAGGTGAACGCACCCATTAAGTGCCTGTGCCTGCACCCCAACCAG
GCAGAGCTCATCGTGGGTGACCAGAGCGGGGCTATCCACATCTGGGACTTGAAAACAGACCACAACGAGC
AGCTGATCCCTGAGCCCGAGGTCTCCATCAGTCCGCCACATCGATCCCGACCCAGCTACATGGCAGC
TGTCATAGCACCAGAACTGCTATGTCTGGAATCTGACGGGGGCGATTGGTGACGAGGTGACCCAGCTC
ATCCCAAGACTAAGATCCCTGCCACACGCGCTACGCCCTGCAGTGTGCTTACGCCCCGACTCCACGC
TCCTCGCCACCTGCTCGGCTGATCAGACGTGCAAGATCTGGAGGACGTCCAATTCTCCCTGATGACGGA
GCTGAGCATCAAGAGCGGCAACCCCGGGAGTCTCCCGCGGCTGGATGTGGGGCTGCGCCTTCTCGGG
GACTCCCAGTACATCGTCACTGCTTCTCGGACAACTGGCCCGGCTCTGGTGTGTGGAGACTGGAGAGA
TCAAGAGAGAGTATGGCGGCCACCAGAAGGCTGTTGTCTGCCTGGCCTTCAATGACAGTGTGCTGGGCTA
G
```



[View online »](#)

5' Read Nucleotide Sequence:

>OriGene 5' read for NM_022372 unedited
 TTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGGCACGAGGGCCGCCCGTAAGAT
 GCTCTGACCTTTGACCCTGCCGTTACAGCTCTAGGGCCCGTGCAGGCCACACCATGAACAC
 CTCCCCAGGCACGGTGGGCAGTGACCCGGTTCATCTGGCCACTGCAGGCTACGACCACAC
 CGTGCGCTTCTGGCAGGCCACAGCGGCATCTGCACCCGGACGGTGCAGCACCAGGACTC
 CCAGGTGAATGCCTTGGAGGTCACACCCGACCGCAGCATGATTGCTGCTGCAGGTTACCA
 GCACATCCGCATGTATGATCTCAACTCCAATAACCCCTAACCCCATCATCAGCTACGACGG
 CGTCAACAAGAACATCGGTCTGTGGGCTTCCACGAAGACGGCCGCTGGATGTACACGGG
 CGGCGAGGACTGCACAGCCAGGATCTGGGACCTCAGGTCCCGAACCTGCAGTGCCAGCG
 GATCTTCCAGGTGAACGCACCCATTAAGTGCCTGCCTGCACCCGAACAGGCAGAGCT
 CATCGTGGGTGACCAGAGCGNGCTATCCACATCTGGGACTTGAAAACAGACCACAACGA
 GCAGCTGATCCCTGAGCCCGAGGCTCCATCACGTCCGCCACATCGATCCCGACGCCAG
 CTACATGGCAGCTGTCAATAGCACCGAAACTGCTATGTCTGGAATCTGNACGGGGGCAT
 TGGTGACGAGGTGACCCAGCTCATCCCAAGACTAAGATCCCTGCCACACGCGCTACGC
 CCTGCAGTGTCTTNCAGCCCCGACTTCACGCTCCTCGCCACCTGCTNGGGTGATCAGAC
 GTGCCAGATCTGNNAGACGTCCAACCTCCTCCCTGATGACGGAGCTGAGCATCAAGAGCG
 GCACCCCGGG

3' Read Nucleotide Sequence:

>OriGene 3' read for NM_022372 unedited
 ACGTAAAAAGAGTGCTTTTCACTTTCCAGATGCGTCCCCCGGACCTGGCCAGCCT
 GCCCGTCCCCTTTCCATGGGCTGGCCAGGCCACCTGCACCAGGGAGGACAGCTGCTGGC
 AGGGACTAATAAACCTTCCACCTGGCCATGGTGGTGGTGTCTCTATGGACCGAGGCC
 TGAAACCGGGCAGGGAGGGGCAGAGAACGCACTAGCTTGGGGTGGGCACCAGCCTCAG
 ACCCTCAGCAGCTTTGGGCCCTCGGCCGACTTCCAGGCAGTGACAGGCTAGCCAGTC
 CAGGAGTGTGCAGCCTGGCTTGGGTCGAGCTCTGTACATCTGGATAAGCAACTGGGGGC
 TGAGAGTCCCAGGGCAGCCTGGCCAGCAGGTCAAGGCGCCACAGGGGGCCATCAGGTCC
 AGCTGGCGCATGCCGGCAGGGGAGGGTCTGCTCTGACCTGAGTGTGCATGGGTCCCTCC
 AGCTGCCACCACCTGCACCAGGCAGTCCCGAGGGGTACAGGCTAGCCCAGCACACTGTC
 ATTGAAGGCCAGGCAGACAACAGCCTTCTGGTGGCCGCCATACTCTCTTGTATCTTCC
 AGTCTCCACACACCAGAGCCGGGCCAGGTTGTCCGAGGAAGCAGTGACGATGTACTIONGA
 GTCCCCCGAGAAGCGCAGCCCCACATTCAGCCCGGGAAGACTTCCCCGGGATTGCCGCT
 CTTGATGCTTACCTCCGTCATAAGGGAGAAGTTGGACGCTCCTCCAGATCTTGCAGTTTTG
 ATCACCCCGACCAGGTGGCCAGGCACCGTGAATCCGGCCTGAACCGACACTGCAGGGCG
 TAACGCGTGTGGCAGGGATTCTATTTTGGCGGATAACCTGGGTACCTGTTACCATGCC
 CCCCTTTAGATCCAACATAGCAGCTTCCGGGCCCTTTTGAACCGCCT

Restriction Sites:

NotI-NotI

ACCN:

NM_022372

Insert Size:

1720 bp

OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_022372.2](#), [NP_071767.2](#)

RefSeq Size: 1931 bp

RefSeq ORF: 984 bp

Locus ID: 64223

UniProt ID: [Q9BVC4](#)

Cytogenetics: 16p13.3

Domains: WD40

Protein Pathways: mTOR signaling pathway

Gene Summary: Subunit of both mTORC1 and mTORC2, which regulates cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Within mTORC1, LST8 interacts directly with MTOR and enhances its kinase activity. In nutrient-poor conditions, stabilizes the MTOR-RPTOR interaction and favors RPTOR-mediated inhibition of MTOR activity. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) represents the longest transcript. Variants 1, 2, and 3 encode the longer isoform (a).