

Product datasheet for SC127424

RPL13 (NM_033251) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RPL13 (NM_033251) Human Untagged Clone

Tag: Tag Free Symbol: RPL13

Synonyms: BBC1; D16S44E; D16S444E; L13; SEMDIST

Mammalian Cell None

Selection:

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene ORF within SC127424 sequence for NM_033251 edited (data generated by NextGen

Sequencing)

GAAGCCGCAGAACAGGATGTTGAAAAGAAAAAATAA

Clone variation with respect to NM $_$ 033251.2

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



5' Read Nucleotide Sequence:

>OriGene 5' read for NM_033251 unedited

TTTTGTATACGACNACTATAGGGCGGCCGCGATTCGGCACGAGGTCCTGCGCAGGAGCCG
CAGGGCCGTAGGCAGCCATGGCGCCCAGCCGGATTGGCATGGTCTTGAAGCCCCACTTCC
ACAAGGACTGGCAGCGGCGCGTGGCCACGTGGTTCAACCAGCCGGCCCGTAAGATCCGCA
GACGTAAGGCCCGGCAAGCCAAGGCGCGCCGCATCGCCCCGCGCCCCGCGTCGGGTCCCA
TCCGGCCCATCGTGCGCTGCCCCACGGTTCGGTACCACACGAAGGTGCGCGCCGCGC
GCTTCAGCCTGGAGGAGCTCAGGGTGGCCGGCATTCACAAGAAGGTGGCCCGGACCATCG
GCATTTCTGTGGATCCGAGGAGGCGGAACAAGTCCACGGAGTCCCTGCAGGCCAACGTGC
AGCGGCTGAAGGAGTACCGCTCCAAACTCATCCTCTTCCCCAGGAAGCCCTCGGCCCCCA
AGAAGGGAGACAGTTCTGCTGAAGAACTGAAACTGGCCACCAGCTGACCGGACCGTCA
TGCCCGTCCGGAACGTCTATAAGAAGGAGAAAACTCGACCCAGCTCATCACTGAGGAAGAAGAA
ATTTCAAAGCCTTCGCTAGTCTCCGTATGGCCCGTGCCAACGCCCGGCTCTTCGGCATAC
GGGCAAAAAGAGCCAAGGAAGCCGCAGAACAGGATGTTGAAAAGAAAAATAAAGCCCTCC
TGGGGACTTGGAATCAGTCGGCAGTCATGCTGGGTCTCCACGTGGTGTTTTCGTGGGAA
CAACTGGCCTGGGATGGGGCTCCACTGCTGTGACTTCCCTGCCAGGGATTTTGGGGCT
TTTTTGAAAGACATCCAAGCCCTGNAT

Restriction Sites: Notl-Notl **ACCN:** NM 033251

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 033251.1, NP 150254.1

 RefSeq Size:
 1296 bp

 RefSeq ORF:
 636 bp

 Locus ID:
 6137

 UniProt ID:
 P26373

 Cytogenetics:
 17p11.2

Domains: Ribosomal_L13e

Protein Families: Druggable Genome

Protein Pathways: Ribosome



Gene Summary:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L13E family of ribosomal proteins. It is located in the cytoplasm. This gene is expressed at significantly higher levels in benign breast lesions than in breast carcinomas. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. [provided by RefSeq, Jul 2011]

Transcript Variant: This variant (2) has a different 5' UTR, compared to variant 1. Variants 1 and 2 encode the same isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.