

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC127643

Dysadherin (FXYD5) (NM_144779) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Dysadherin (FXYD5) (NM_144779) Human Untagged Clone
Tag:	Tag Free
Symbol:	Dysadherin
Synonyms:	DYSAD; HSPC113; IWU1; KCT1; OIT2; PRO6241; RIC
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC127643 representing NM_144779. Blue=Insert sequence <mark>Red</mark> =Cloning site Green=Tag(s)
	GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC ATGTCGCCCTCTGGTCGCCTGTGTCTTCTCACCATCGTTGGCCTGATTCTCCCCACCAGAGGACAGACG TTGAAAGATACCACGTCCAGTTCTTCAGCAGACTCAACTATCATGGACATTCAGGTCCCGACACGAGCC CCAGATGCAGTCTACACAGAACTCCAGCCCACCTCTCCAACCCAACCTGGCCTGCTGATGAAACACCA CAACCCCAGACCCAGACCCAGCAACTGGAAGGAACGGATGGGCCTCTAGTGACAGATCCAGAGACACAC AAGAGCACCAAAGCAGCTCATCCCACTGATGACACCACGACGCTCTCTGAGAGAACCATCCCAGAGACACAC GACGTCCAGACAGACCCCCAGACCTCAAGCCATCTGGTTTTCATGAGGGATGACCCCTTCTTCTATGAT GAACACACCCTCCGGAAACGGGGGCTGTTGGTCGCCGCTGTGGCTGTTCATCACAGGCATCATCCTCC ACCAGTGGCAAGTGCAGGCAGCTGTTCGTCGCGGTATGCCGGAATCGTTGCAGGTGA ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
Chromatograms:	https://cdn.origene.com/chromatograms/mk8074_c09.zip
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_144779
Insert Size:	537 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Dysadherin (FXYD5) (NM_144779) Human Untagged Clone – SC127643
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution N	 thod: 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 144779.1</u>
RefSeq Size:	927 bp
RefSeq ORF:	537 bp
Locus ID:	53827
UniProt ID:	<u>Q96DB9</u>
Cytogenetics:	19q13.12
Protein Families	Druggable Genome, Ion Channels: Other, Transmembrane
MW:	19.5 kDa
Gene Summary:	This gene encodes a member of a family of small membrane proteins that share a 35-amino acid signature sequence domain, beginning with the sequence PFXYD and containing 7 invariant and 6 highly conserved amino acids. The approved human gene nomenclature for the family is FXYD-domain containing ion transport regulator. Mouse FXYD5 has been termed RIC (Related to Ion Channel). FXYD2, also known as the gamma subunit of the Na,K-ATPase, regulates the properties of that enzyme. FXYD1 (phospholemman), FXYD2 (gamma), FXYD3 (MAT-8), FXYD4 (CHIF), and FXYD5 (RIC) have been shown to induce channel activity in experimental expression systems. Transmembrane topology has been established for two family members (FXYD1 and FXYD2), with the N-terminus extracellular and the C-terminus on the cytoplasmic side of the membrane. This gene product, FXYD5, is a glycoprotein that functions in the up-regulation of chemokine production, and it is involved in the reduction of cell adhesion via its ability to down-regulate E-cadherin. It also promotes metastasis, and has been linked to a variety of cancers. Alternative splicing results in multiple transcript variants. [RefSeq curation by Kathleen J. Sweadner, Ph.D., sweadner@helix.mgh.harvard.edu., Sep 2009] Transcript Variant: This variant (1) encodes the same protein as variants 2 and 3.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US