

Product datasheet for **SC200153**

RPL5 (NM_000969) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	RPL5 (NM_000969) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	RPL5
Synonyms:	L5; MSTP030; PPP1R135; uL18
ACCN:	NM_000969
Insert Size:	89 bp
Insert Sequence:	>SC200153 3'UTR clone of NM_000969 The sequence shown below is from the reference sequence of NM_000969. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC AGAGCTCAGGAGCGGGCTGCTGAGAGCTAAACCCAGCAATTTTCTATGATTTTTTCAGATATAGATAAT AAACTTATGAACAGCAACTA ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM_000969.5</u>



[View online »](#)

Summary:

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of four RNA species and approximately 80 structurally distinct proteins. This gene encodes a member of the L18P family of ribosomal proteins and component of the 60S subunit. The encoded protein binds 5S rRNA to form a stable complex called the 5S ribonucleoprotein particle (RNP), which is necessary for the transport of nonribosome-associated cytoplasmic 5S rRNA to the nucleolus for assembly into ribosomes. The encoded protein may also function to inhibit tumorigenesis through the activation of downstream tumor suppressors and the downregulation of oncoprotein expression. Mutations in this gene have been identified in patients with Diamond-Blackfan Anemia (DBA). This gene is co-transcribed with the small nucleolar RNA gene U21, which is located in its fifth intron. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed throughout the genome. [provided by RefSeq, Mar 2017]

Locus ID:

6125

MW:

3.4