

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC200239

HIST1H2AK (HIST1H2AG) (NM_021064) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	HIST1H2AK (HIST1H2AG) (NM_021064) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	H2AC11
Synonyms:	H2A.1b; H2A/p; H2AC13; H2AC15; H2AC16; H2AC17; H2AFP; H2AG; HIST1H2AG; pH2A/f
ACCN:	NM_021064
Insert Size:	101 bp
Insert Sequence:	>SC200239 3'UTR clone of NM_021064 The sequence shown below is from the reference sequence of NM_021064. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site
	GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAA <mark>GCGATCGC</mark> GAGAGCCACCACAAGGCGAAGGGCAAGTAACTATCTGTACTAGTTTGTGGCAGCTCAAGTAAAATCGAG TCCAAACCAACGGCTCTTTTCAGGGCCACCCA ACGCGTAAGCGGCCGCGGCATCTAGATTCGAAGAAAATGACCGACC
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM 021064.5</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	HIST1H2AK (HIST1H2AG) (NM_021064) Human 3' UTR Clone – SC200239
Summary:	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015]
Locus ID:	8969
MW:	3.5

~_____

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US