

Product datasheet for **SC200327**

ATP5PD (NM_001003785) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	ATP5PD (NM_001003785) Human 3' UTR Clone
Symbol:	ATP5PD
Synonyms:	APT5H; ATP5H; ATPQ
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_001003785
Insert Size:	99 bp
Insert Sequence:	<p>>SC200327 3'UTR clone of NM_001003785</p> <p>The sequence shown below is from the reference sequence of NM_001003785. The complete sequence of this clone may contain minor differences, such as SNPs.</p> <p>Blue=Stop Codon Red=Cloning site</p> <pre> GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGCCGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC TGGCCTCACCAACCAATTGAGAATTTAAAAAATTGAGTCCAGGAGGAAGCTCTGGCCCTTGATTACAC ATTCTGGACATTAAAAATAATAATTATACA ACGCGTAAGCGGCCGCGCATCTAGATTCTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG </pre>
Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM_001003785.2</u>


[View online »](#)

Summary:

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The F1 complex consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled in a ratio of 3 alpha, 3 beta, and a single representative of the other 3. The Fo seems to have nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the d subunit of the Fo complex. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. In addition, three pseudogenes are located on chromosomes 9, 12 and 15. [provided by RefSeq, Jun 2010]

Locus ID:

10476

MW:

3.9