

## **Product datasheet for SC200344**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

## DMAP1 (NM 001034023) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: DMAP1 (NM\_001034023) Human 3' UTR Clone

**Vector:** pMirTarget (PS100062)

Symbol: DMAP1

Synonyms: DNMAP1; DNMTAP1; EAF2; MEAF2; SWC4

**ACCN:** NM\_001034023

**Insert Size:** 85 bp

Insert Sequence: >SC200344 3'UTR clone of NM\_001034023

The sequence shown below is from the reference sequence of NM\_001034023. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

TCTTCCGTGAAGAAGCCAAGAAGCCGTGAGAGGCCCCACGGGGTGTGGGCCGACGCTGTTATGTAAATA

GAGCTGCTGAGTTGGA

 ${\tt CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG}$ 

Restriction Sites: Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** NM 001034023.2





ORIGENE

**Summary:** 

This gene encodes a subunit of several, distinct complexes involved in the repression or activation of transcription. The encoded protein can independently repress transcription and is targeted to replication foci throughout S phase by interacting directly with the N-terminus of DNA methyltransferase 1. During late S phase, histone deacetylase 2 is added to this complex, providing a means to deacetylate histones in transcriptionally inactive heterochromatin following replication. The encoded protein is also a component of the nucleosome acetyltransferase of H4 complex and interacts with the transcriptional corepressor tumor susceptibility gene 101 and the pro-apoptotic death-associated protein 6, among others. Alternatively spliced transcript variants encoding the same protein have been described. [provided by RefSeq, Jul 2008]

**Locus ID:** 55929

**MW:** 3