

Product datasheet for SC200516

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

H2BC5 (NM_021063) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: H2BC5 (NM_021063) Human 3' UTR Clone

Symbol: H2BC5

Synonyms: dJ221C16.6; H2B.1B; H2B/a; H2B/b; H2B/g; H2B/h; H2B/k; H2B/k; H2BFA; H2BFB; H2BFG;

H2BFH; H2BFK; H2BFL; HIRIP2; HIST1H2BC; HIST1H2BD; HIST1H2BE; HIST1H2BF; HIST1H2BG;

HIST1H2BI

Mammalian Cell

Selection:

Neomycin

Vector: pMirTarget (PS100062)

ACCN: NM_021063

Insert Size: 87 bp

Insert Sequence: >SC200516 3'UTR clone of NM_021063

The sequence shown below is from the reference sequence of NM_021063. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GCCGTCACCAAGTACACCAGTTCCAAGTAACTTTGCCAAGTAAGCATCTTTACACCTAATCCCAAAGGC

TCTTTTAAGAGCCACGCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 021063.4</u>





ORÏGENE

MW:

Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2B family. Two transcripts that encode the same protein have been identified for this gene, which is found in the large histone gene cluster on chromosome

6p22-p21.3. [provided by RefSeq, Aug 2015]

Locus ID: 3017

3.1

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US